

Original Publications

- (295) Yoko Irukayama-Tomobe, Yasuhiro Ogawa, Hiromu Tominaga, Yukiko Ishikawa, Naoto Hosokawa, Yuki Kawabe, Shuntaro Uchida, Shinobu Ambai, Ryo Nakajima, Tsuyoshi Saito, Takeshi Sakusai, Hiroshi Nagase, and Masashi Yanagisawa, A non-peptide orexin type 2 receptor agonist ameliorates narcolepsy-cataplexy symptom in mouse model, *PNAS*, *114* (22), 5731-5736, **2017**. IF: 9.423,
- (294) Kohei Hayashida, Shigeto Hirayama, Takashi Iwai, Yoshikazu Watanabe, Toshihiro Takahashi, Junichi Sakai, Eriko Nakata, Tomio Yamakawa, Hideaki Fujii, Hiroshi Nagase, Novel delta opioid receptor agonists with oxazatricyclodecane structure 2, *Bioorg. Med. Chem. Lett.*, *27*, 2742-2745, **2017**, IF: 2.486
- (293) Katherine Lansu, Joel Karpiak, Jing Liu, Xi-Ping Huang, John D McCorvy, Wesley K Kroeze, Tao Che, Hiroshi Nagase, Frank I Carroll, Jian Jin, Brian K Shoichet & Bryan L Roth, In silico design of novel probes for the atypical opioid receptor MRGPRX2, *Nature Chemical Biology*. *13* (5), 529-536, **2017** IF:12.709,
- (292) Yusuke Hamada, Yoshihiro Tasaki, Kana Morita, Kohei Yamamizu, Michiko Narita, Fukiko Matuyama, Masami Suzuki, Daigo Ikegami, Kazuhiro, Arakawa, Yasuyuki Nagumo, Miho Kawata, Yasuhiro Uezono, Hiroshi Nagase, Kazunori, Aoki, Jun Yamashita, Noako Kuzumaki, Minoru Narita, Nalfurafine enhances chemotherapy-induced survival advantage, *Jpn. J. Pharm.Palliat. Care Sci.* *10*, 7-12, **2017**.
- (291) Hiroshi Nagase, Naoshi Yamamoto, Masahiro Yata, Sayaka Ohruai, Takahiro Okada, Tsuyoshi Saitoh, Noriki Kutsumura, Yasuyuki Nagumo, Yoko Irukayama-Tomobe, Yukiko Ishikawa, Yasuhiro Ogawa, Shigeto Hirayama, Daisuke Kuroda, Yurie Watanabe, Hiroaki Gouda, and Masashi Yanagisawa, Design and Synthesis of Potent and Highly Selective Orexin 1 Receptor Antagonists with a Morphinan Skeleton and their Pharmacologies, *J. Med. Chem.*, *60*, 1018-1040, **2017**.
- (290) Leo Gotoh, Akiyoshi Saitoh, Misa Yamada, Hideaki Fujii, Hiroshi Nagase, Mitsuhiko Yamada, Effects of repeated treatment with a delta opioid receptor agonist KNT-127 on hyperemotionality in olfactory-bulbectomized rats, *Behav. Brain Res.*, *323*, 11-14 **2017**.
- (289) Saki Yajima, Tsuyoshi Saitoh, Kohei Kawa, Kensuke Nakamura, Hiroshi Nagase, Yasuaki Einaga, Shigeru Nishiyama, Asymmetric induction in

- cyclohexadienones carrying α -D- glucopyranosyl Moiety, *Tetrahedron*, 51, 8428-8435, **2016**.
- (288) Noriki Kutsumura, Yasuaki Koyama, Kotaro Tateno, Naoshi Yamamoto, Hiroshi Nagase, Takao Saito, ZnCl₂-Promoted Intramolecular Hetero Diels-Alder Reaction of *o*-Alkynylphenylcarbodiimides for Synthesis of Dihydrodibenzo[b,g][1,8]naphthyridines, *Chem. Pharm. Bull.*, 64, 1364-1369, **2016**.
- (287) Chrismawan Ardianto, Naomi Yonemochi, Shogo Yamamoto, Lizhe Yang, Fumiko Takenoya, Seiji Shioda, Hiroshi Nagase, Hiroko Ikeda, Junzo Kamei, Opioid System in the Lateral Hypothalamus Regulate Feeding Behavior Through Orexin and GABA Neurons, *Neuroscience*, 320, 183-193, **2016**.
- (286) Takaaki Komatsu, Soh Katsuyama, Hiroshi Nagase, Hirokazu Mizoguchi, Chikai Sakurada, Minoru Tsuzuki, Shinobu Sakurada, Tsukasa Sakurada, Intrathecal morphine-3-glucuronide-induced nociceptive behavior via Delta-2 opioid receptors in the spinal cord, *Pharmacology Biochemistry and Behavior* 140, 68-74, **2016**.
- (285) Ryo Nakajima, Naoshi Yamamoto, Shigeto Hirayama, Takashi Iwai, Akiyoshi Saitoh, Yasuyuki Nagumo, Hideaki Fujii, Hiroshi Nagase, Synthesis of new opioid derivatives with a propellane skeleton and their pharmacologies: Part 5, novel pentacyclic propellane derivatives with a 6-amide side chain, *Bioorg. Med. Chem.*, 23, 6271-6279, **2015**.
- (284) Takashi Nagahara, Tsuyoshi Saitoh, Noriki Kutsumura, Yoko Irukayama, Tomobe, Yasuhiro Ogawa, Daisuke Kuroda, Hiroaki Gouda, Hidetoshi Kumagai, Hideaki Fujii, Masashi Yanagisawa, Hiroshi Nagase, Design and Synthesis of Non-Peptide, Selective Orexin Receptor 2 Agonists, *J. Med. Chem.*, 58, 7931-7937, **2015**.
- (283) Hiroko Ikeda, Naomi Yonemochi, Lizhe Yang, Takahiro Ohashi, Megumi Ikegami, Hiroshi Nagase, Junzo Kamei, Inhibition of opioid systems in the hypothalamus as well as the mesolimbic area suppresses feeding behavior of mice, *Neuroscience*, 311, 9-21, **2015**.
- (282) Noriki Kutsumura, Ryo Nakajima, Yasuaki Koyama, Yoshiyuki Miyata, Tsuyoshi Saitoh, Naoshi Yamamoto, Satoshi Iwata, Hideaki Fujii, Hiroshi Nagase, Investigation of 7-benzylidenenaltrexone derivatives as a novel structural antitrichomonal lead compound, *Bioorg. Med. Chem. Lett.*, 25, 4890-4892, **2015**.
- (281) Toru Nemoto, Yusuke Iihara, Shigeto Hirayama, Takashi Iwai, Eika Higashi,

- Hideaki Fujii, Hiroshi Nagase, Naltrindole derivatives with fluorinated ethyl substituents on 17-nitrogen as δ opioid receptor inverse agonist, *Bioorg. Med. Chem. Lett.*, 2927-2930, *25*, **2015**. IF:2.427
- (280) Masami Suzuki, Fumiko Chiwaki, Yumi Sawada, Maho Ashikawa, Kazuhiko Aoyagi, Takeshi Fujita, Kazuyoshi Yanagihara, Masayuki Komatsu, Minoru Narita, Tsutomu Suzuki, Hiroshi Nagase, Ryoji Kushima, Hiromi Sakamoto, Takeo Fukagawa, Hitoshi Katai, Hitoshi Nakagama, Teruhiko Yoshida, Yasuhito Uezono, Hiroki Sasaki, Peripheral Opioid Antagonist Enhances the Effect of Anti-Tumor Drug by Blocking a Cell Growth-Suppressive Pathway In Vivo, *PLoS One*, 10, e0123407, **2015**.
- (279) Kazuya Konoura, Hideaki Fujii, Satomi Imaide, Hiroaki Gouda, Shigeto Hirayama, Shuichi Hirono, Hiroshi Nagase, Transformation of naltrexone into mesembrane and investigation of the binding properties of its intermediate derivatives to opioid receptors, *Bioorg. Med. Chem.*, 23, 439-448, **2015**.
- (278) Shigeto Hirayama, Naohisa Wada, Naoya Kuroda, Takashi Iwai, Noriyuki Yamaotsu, Shuichi Hirono, Hideaki Fujii, Hiroshi Nagase, Synthesis of a novel universal opioid receptor agonist with the 1,3,5-trioxazatriquinane skeleton and its pharmacologies, *Bioorg. Med. Chem. Lett.*, 24, 4895-4898, **2014**.
- (277) Chihiro Nozaki, Hiroshi Nagase, Toru Nemoto, Audrey Matifas, Claire Gaveriaux-Ruff, and Brigitte L. Kieffer, *In vivo* properties of KNT-127, a novel delta opioid agonist: receptor internalization, antihyperalgesia and antidepressant effects in mice, *Br. J. Pharmacol.*, 171, 5376-5386, **2014**.
- (276) Yoshikazu Watanabe, Shota Kitazawa, Hideaki Fujii, Toru Nemoto, Shigeto Hirayama, Takashi Iwai, Hiroaki Gouda, Shuichi Hirono, Hiroshi Nagase, Design, synthesis, and structure-activity relationship of novel opioid κ receptor selective agonists: α -Iminoamide derivatives with an azabicyclo[2.2.2]octene skeleton, *Bioorg. Med. Chem. Lett.*, 24, 4980-4983, **2014**.
- (275) Shigeto Hirayama, Naohisa Wada, Toru Nemoto, Takashi Iwai, Hideaki Fujii, Hiroshi Nagase, Synthesis and Pharmacology of a Novel κ Opioid Receptor (KOR) Agonist with a 1,3,5-Trioxazatriquinane Skeleton, *ACS Med. Chem. Lett.*, 5, 868-872, **2014**.
- (274) Azusa Sugiyama, Hiroshi Nagase, Jun-ichiro Oka, Mitsuhiro Yamada, Akiyoshi Saito, DOR2-selective but not DOR1-selective antagonist abolishes anxiolytic-like effects of the δ opioid receptor agonist KNT-127, *Neurobiology*, 11, 314-320, **2014**.
- (273) Hiroshi Nagase, Ryo Nakajima, Naoshi Yamamoto, Shigeto Hirayama, Takashi

- Iwai, Toru Nemoto, Hiroaki Gouda, Shuichi Hirono, Hideaki Fujii, Design and Synthesis of quinolinopropellane Derivatives with Selective Delta Opioid Receptor Agonism, *Bioorg. Med. Chem. Lett.*, 24, 2851-2854, **2014**.
- (272) Hideaki Fujii, Kohei Hayashida, Akiyoshi Saitoh, Akinobu Yokoyama, Shigeto Hirayama, Takashi Iwai, Eriko Nakata, Toru Nemoto, Yuka Sudo, Yasuhito Uezono, Mitsuhiko Yamada, Hiroshi Nagase, Novel Delta Opioid Receptor Agonists with Oxazatricyclodecane Structure, *ACS Med. Chem. Lett.*, 5, 368-372, **2014**.
- (271) Adam Kardon, Erika Polgar, Junichi Hachisuka, Darren Cameron, Sinead Savage, Lindsey McClement, Xiaoyun Cai, Sergei Karnup, Christopher Fan, Gregory Hemenway, Carcha Bernard, Erica S.Schwartz, Hiroshi Nagase, Masahiko Watanabe, Takahiro Furuta, Takeshi Kaneko, Andrew J. Todd, Sarah E. Ross, Dynorphin acts as a neuromodulator to inhibit itch in the dorsal horn of the spinal cord, *Neuron*. 82, 573-586, **2014**.
- (270) Kyoko Ishikawa, Miyuki Tomatsu, Hiroshi Nagase, and Hideaki Fujii, Zinc-acetic acid promoted reductive carbon-nitrogen bond cleavage reaction of α -aminoketones, *Heterocycles*, 88, 1051-1063, **2014**.
- (269) Kohei Yamamizu, Sadayoshi Furuta, Yusuke Hamada, Akira Yamashita, Naoko Kuzumaki, Michiko Narita, Kento Doi, Shiori Katayama, Hiroshi Nagase, Jun Yamashita, Minoru Narita, κ Opioids inhibit tumor angiogenesis by suppressing VEGF signaling, *Scientific report*, 3, 3213, **2013**.
- (268) Toru Nemoto, Yoshihiro Ida, Yusuke Iihara, Ryo Nakajima, Shigeto Hirayama, Takashi Iwai, Hideaki Fujii, Hiroshi Nagase, The most effective influence of 17-(3-ethoxypropyl) substituent on the binding affinity and the agonistic activity in KNT-127 derivatives, δ opioid receptor agonists, *Bioorg. Med. Chem.*, 21, 7628-7647, **2013**.
- (267) Yoshikazu Watanabe, Shota Kitazawa, Toru Nemoto, Shigeto Hirayama, Takashi Iwai, Hideaki Fujii, Hiroshi Nagase, Design and synthesis of novel opioid ligands with an azabicyclo[2.2.2]octane skeleton having a 7-amide chain and their pharmacologies, *Bioorg. Med. Chem.*, 21, 3032-3050, **2013**.
- (266) Yuya Ise, Tomohisa Mori, Shirou Katayama, Hiroshi Nagase, Tsutomu Suzuki, Rewarding effects of ethanol combined with low doses of morphine through dopamine D1 receptor, *J. Nippon Medical School*, 80, No.1, 34-41, **2013**.
- (265) Akiyoshi Saitoh, Azusa Sugiyama, Misa Yamada, Masatoshi Inagaki, Jun-Ichiro Oka, Hiroshi Nagase, Mitsuhiko Yamada, The novel δ opioid receptor agonist KNT-127 produces distinct anxiolytic-like effect in rats without producing the

- adverse effects associated with benzodiazepines, *Neuropharmacology*, *67*, 485-493, **2013**.
- (264) Toru Nemoto, Shigeto Hirayama, Takashi Iwai, Naoshi Yamamoto, Yukimasa Harada, Naohisa Wada, Miyuki Tomatsu, Marina Ishihara, Hideaki Fujii, Hiroshi Nagase, The effect of 17-N-substituents on the activity of the opioid κ receptor in nalfurafine derivatives, *Bioorg. Med. Chem. Lett*, *23*, 268-272, **2013**.
- (263) Enomoto, T.; Yamashita, A.; Torigoe, K.; Horiuchi, H.; Hirayama, S.; Nakahara, K.; Yanase, M.; Sakai, H.; Ikegami, D.; Nagase, H.; Suzuki, T.; Iseki, M.; Inada, E.; Narita, M. **Effects of mirtazapine on sleep disturbance under neuropathic pain-like state**, *Synapse*, *66*, 483-488, **2012**. IF: 2.945
- (262) Hideaki Fujii, Takaaki Yamada, Kohei Hayashida, Miki Kuwada, Atom Hamasaki, Kazunori, Nobuhara, Sumio Ozeki, and Hiroshi Nagase, Application of SO₃H Silica gel to Deprotection of Silyl Ethers, *Heterocycles*, *85*, No. 11, 2685-2691, **2012**.
- (261) Hideaki Fujii, Ryo Nakajima, Junko Akiyama, Naoshi Yamamoto, Shigeto Hirayama, Toru Nemoto, Hiroaki Gouda, Shuichi Hirono, Hiroshi Nagase, Synthesis of new opioid derivatives with a propellane skeleton and their pharmacology: Part 3, novel propellane derivatives with pentacyclic skeletons, *Bioorg. Med. Chem. Lett.*, *22*, 7697-7701, **2012**.
- (260) Yoshihiro Ida, Ayaka Matsubara, Toru Nemoto, Manabu Saito, Shigeto Hirayama, Hideaki Fujii, Hiroshi Nagase, Synthesis of quinolinomorphinan derivatives as highly selective δ opioid receptor ligands, *Bioorg. Med. Chem.*, *20*, 5810-5831, **2012**.
- (259) Hideaki Fujii, Hiroshi Nagase, Essential structure of opioid κ receptor agonist nalfurafine for binding to the κ receptor 3: Synthesis of decahydroiminoethano)phenanthren derivatives with an oxygen functionality at the 3-position and their pharmacologies, *Bioorg. Med. Chem. Lett.*, *22*, 7711-7714, **2012**.
- (258) Naohisa Wada, Hideaki Fujii, Koji Koyano, Shigeto Hirayama, Takashi Iwai, Toru Nemoto, and Hiroshi Nagase, Synthesis of novel triplet drugs with 1,3,5-trioxatriquinane skeletons and their pharmacologies. 3: Synthesis of novel triplet drugs with the bis(epoxymethano) or bis(dimethylepoxymethano) structure (double-capped triplet), *Bioorg. Med., Chem., Lett.*, *22*, 7551-7554, **2012**.
- (257) Yoshinori Miyata, Hideaki Fujii, Yuka Uenohara, Seiki Kobayashi, Tsutomu

- Takeuchi, Hiroshi Nagase, Investigation of 7-benzylidenenaltrexone derivatives as reverser of resistance for chloroquine-resistant *Plasmodium chabaudi*, *Bioorg. Med. Chem. Lett.*, *22*, 5174-5176, **2012**.
- (256) Hiroshi Nagase, Satomi Imaide, Shigeto Hirayama, Toru Nemoto, Hideaki Fujii, Essential structure of opioid κ receptor agonist nalfurafine for binding to the κ receptor 2: Synthesis of decahydroiminoethano)phenanthrene derivatives and their pharmacologies, *Bioorg. Med. Chem. Lett.*, *22*, 5071-5074, **2012**.
- (255) Hiroshi Nagase, Satomi Imaide, Takaaki Yamada, Shigeto, Hirayama, Toru Nemoto, Noriyuki Yamaotsu, Shuichi Hirono, and Hideaki Fujii, Essential Structure of Opioid κ Receptor Agonist Nalfurafine for Binding to κ Receptor 1: Synthesis of Decahydroisoquinoline Derivatives and Their Pharmacologies, *Chem. Pharm., Bull.*, *60(8)*945-948, **2012**.
- (254) Yoshikazu Watanabe, Shota Kitazawa, Hideaki Fujii, Toru Nemoto, Shigeto Hirayama, Hiroshi Nagase, Design and synthesis of novel opioid ligands with an azabicyclo[2.2.2]octane skeleton and their pharmacologies, *Bioorg. Med. Chem.*, *22*, 2689-2692, **2012**.
- (253) Yoshihiro Ida, Toru, Nemoto, Shigeto Hirayama, Hideaki Fujii, Yumiko Osa, Masayuki Imai, Takashi Nakamura, Toshiyuki Kanemasa, Akira Kato, Hiroshi Nagase, Synthesis of quinolinomorphinan-4-ol derivatives as delta opioid receptor agonist, *Bioorg. Med. Chem.* *20*, 949-961, **2012**.
- (252) Kazuhiro Torigoe, Kae Nakahara, Mahardian Rahmadi, Kazumi Yoshizawa, Hiroshi Horiuchi, Shigeto Hirayama, Satoshi Imai, Naoko Kuzumaki, Toshimasa Ito, Akira Yamashita, Kiyoshi Shakunaga, Mitsuaki Yamasaki, Hiroshi Nagase, Motohiro Matoba, Tsutomu Suzuki, Minoru Narita, Usefulness of Olanzapine as an Adjunct to Opioid Treatment and for the Treatment of Neuropathic pain, *Anesthesiology*, *116*, 159-169, **2012**.
- (251) Shunsuke Tanahashi, Yuko Ueda, Satoshi Yamamura, Hiroshi Nagase, Motohiro Okada, Novel δ 1-receptor agonist KNT-127 increases releases of dopamine and L-Glutamate in striatum, nucleus accumbens and median prefrontal cortex, *Neurpharmacology*, *62*, 2057-2067, **2012**.
- (250) Hiroshi Nagase, Junko Akiyama, Ryo Nakajima, Shigeto Hirayama, Toru Nemoto, Hiroaki Gouda, Shuichi Hirono, Hideaki Fujii, Synthesis of new opioid derivatives with a propellane skeleton and their pharmacology: Part 2, propellane derivatives with an amide side chain *Bioorg. Med. Chem. Lett.*, *22*, 2775-2779, **2012**.
- (249) Naoko Kuzumaki, Atsuo Suzuki, Michiko Narita, Takahiro Hosoya, Atsumi

- Nagasawa, Satoshi Imai, Kohei Yamamizu, Hiroshi Morita, Hiroshi Nagase, Yohei Okada, Hirotaka James Okano, Jun K Yamashita, Hideyuki Okano, Tsutomu Suzuki, and Minoru Narita, Effect of κ -opioid, receptor agonist on the growth of non-small-cell lung cancer (NSCLC) cells, *British J. Cancer*, 106, (6), 1148-1152, **2012**,
- (248) Tatsuya Enomoto, Akira Yamashita, Kazuhiro Torigoe, Hiroshi Horiuchi, Shigeto Hirayama, Kae Nakahara, Makoto Yanase, Hiroyasu Sakai, Ikegami, Daigo, Hiroshi Nagase, Tsutomu Suzuki, Masako Iseki, Eiichi Inada, Minoru Narita, Effects of mirtazapine on sleep disturbance under neuropathic pain-like state, *Synapse*, Dec 28, doi:10.1002/syn.21532, **2011**.
- (247) Yoshinori Miyata, Hideaki Fujii, Yumiko Osa, Seiki Kobayashi, Tsutomu Takeuchi, Hiroshi Nagase, Opioid δ_1 receptor antagonist 7-benzylidenenaltrexone as an effective resistance reverser for chloroquine-resistant *Plasmodium chabaudi*, *Bioorg. Med. Chem. Lett.*, 21, 4710-4712, **2011**.
- (246) Hiroshi Nagase, Koji Koyano, Naohisa Wada, Shigeto Hirayama, Akio Watanabe, Toru Nemoto, Mayumi Nakajima, Koaru Nakao, Hidenori Mochizuki, Hideaki Fujii, Synthesis of novel triplet drugs with 1,3,5-trioxazatriquinane skeletons and their pharmacologies. 2, synthesis of novel triplet drugs with epoxyethano-structure (capped homotriplet), *Bioorg. Med. Chem. Lett.*, 21, 6198-6202, **2011**.
- (245) Hiroshi Nagase, Akio Watanabe, Toru Nemoto, Mayumi Nakajima, Ko Hasebe, Hidenori Mochizuki, and Hideaki Fujii, Synthesis of novel triplet drugs with 1,3,5-trioxazatriquinane skeletons and their pharmacologies. 1: Synthesis of triplet drugs with morphinan skeletons, *Bioorg. Med. Chem. Lett.*, 21, 4023-4026, **2011**.
- (244) Kohei Hayashida, Hideaki Fujii, Shigeto Hirayama, Toru Nemoto, Hiroshi Nagase, Rearrangement of 4,5 - α -Epoymorphinan Derivatives With Carbamoyl epoxy Rings Provide Novel Oxazatricyclodecane Structures, *Tetrahedron*, 67, 6682-6688, **2011**.
- (243) Naoshi Yamamoto, Hideaki Fujii, Toru Nemoto, Ryo Nakajima, Shinobu Momen, Naoki Izumimoto, Ko Hasebe, Hidenori Mochizuki, Hiroshi Nagase, Synthesis of new opioid derivatives with a propellane skeleton and their pharmacology : Part1, *Bioorg. Med. Chem. Lett.*, 21, 4104-4107, **2011**.
- (242) Yamamizu K, Furuta S, Narita M, Kuzumaki N, Imai S, Nagase H, Suzuki T, Narita M, Yamashita JK, *Blood*, **2011**, Jul 21, 118(3); 775-85. Epub **2011**, Apr

- 1.
- (241) Akiyoshi Saito, Azusa Sugiyama, Toru Nemoto, Hideaki Fujii, Kenji Wada, Jun-ichiro Oka, Hiroshi Nagase, Mitsuhiko Yamada, The novel δ opioid receptor agonist KNT-127 produces distinct anxiolytic-like effects in rats without producing the adverse effects associated with benzodiazepines, *Behavioral Brain Res.* 223, 271-279, **2011**.
- (240) Toru Nemoto, Naoshi Yamamoto, Akio Watanabe, Hideaki Fujii, Ko Hasebe, Mayumi Nakajima, Hidenori Mochizuki, Hiroshi Nagase, Synthesis of 6,14-epoxymorphinan derivatives and their pharmacologies, *Bioorg. & Med. Chem.* 19, 1205-1221, **2011**.
- (239) Naoshi Yamamoto, Hideaki Fujii, Satomi Imaide, Toru Nemoto, Shigeto Hirayama, Junji Inokoshi, Hiroshi Tomoda, Hiroshi Nagase, Synthesis of (-)-Homogalanthamine from Naltrexone, *J. Org. Chem.*, 76, 2257-2260, **2011**.
- (238) Hiroshi Nagase, Toru Nemoto, Ayaka Matsubara, Manabu Saito, Naoshi Yamamoto, Yumiko Osa, Shigeto Hirayama, Mayumi Nakajima, Kaoru Nakao, Hidenori Mochizuki, Hideaki Fujii, Design and synthesis of KNT-127, a δ -opioid agonist effective by systemic administration, *Bioorg. & Med. Chem. Lett.*, 20, 6302-6305, **2010**.
- (237) Hideaki Fujii, Yoshihiro Ida, Shinichi Hanamura, Yumiko Osa, Toru Nemoto, Mayumi Nakajima, Ko Hasebe, Kaoru Nakao, Hidenori, Mochizuki, Hiroshi Nagase, Synthesis of pyrrolomorphinan derivatives as κ opioid agonists, *Bioorg. & Med. Chem. Lett.*, 20, 5035-5038, **2010**.
- (236) Noriyuki Yamaotsu, Hideaki Fujii, Hiroshi Nagase, Shuichi Hirono, Identification of the three dimensional pharmacophore of κ -opioid receptor agonists, *Bioorg. & Med. Chem.*, 18, 4446-4452, **2010**.
- (235) Hiroshi Nagase, Satomi Imaide, Miyuki Tomatsu, Shigeto Hirayama, Toru Nemoto, Mayumi Nakajima, Kaoru Nakao, Hidenori Mochizuki, Hiroaki Gouda, Shuichi Hirono, Hideaki Fujii, Investigation of Beckett-Casy model 3: Synthesis of novel naltrexone derivatives with contracted and expanded D-rings and their pharmacology, *Bioorg. & Med. Chem. Lett.*, 20, 3801-3804, **2010**.
- (234) Hiroshi Nagase, Satomi Imaide, Miyuki Tomatsu, Toru Nemoto, Mayumi Nakajima, Kaoru Nakao, Hidenori Mochizuki, Hideaki Fujii, Investigation of Beckett-Casy model 2: Synthesis of novel 15-16nornaltrexone derivatives and their pharmacology *Bioorg. & Med. Chem. Lett.*, 20, 3726-3729, **2010**
- (233) Hideaki Fujii, Imaide Satomi, Akio Watanabe, Kenji Yoza, Mayuki Nakajima, Kaoru Nakao, Hidenori Mochizuki, Noriko Sato, Toru, Nemoto,

- Hiroshi Nagase, A Double Decarboxylation Reaction of an Oxazolidinone and Carboxylic Acid: Its Application to the Synthesis of a New Opioid Lead Compound. *J. Org. Chem.*, *75*, 3, 995-998, **2010**.
- (232) Satomi Imaide, Hideaki Fujii, Akio Watanabe, Toru Nemoto, Mayumi Nakajima, Kaoru Nakao, Hidenori Mochizuki, Hiroshi Nagase, Investigation of Beckett-Casy Model 1: Synthesis of novel 16,17-seco-naltrexone derivatives and their pharmacology. *Bioorg. & Med. Chem. Lett.*, *20*, 1055-1058, **2010**.
- (231) Hiroshi Nagase, Akio Watanabe, Toru Nemoto, Noriyuki Yamaotsu, Kohei Hayashida, Mayumi Nakajima, Ko Hasebe, Kaoru Nakao, Hidenori Mochizuki, Shuichi Hirono, Hideaki Fujii, Drug design and synthesis of a novel kappa opioid receptor agonist with an oxabicyclo[2.2.2]octane skeleton and its pharmacology. *Bioorg. & Med. Chem. Lett.*, *20*, 121-124, **2010**.
- (230) Hideaki Fujii, Kayoko Okada, Marina Ishihara, Shinichi Hanamura, Yumiko Osa, Toru Nemoto; Hiroshi Nagase, Hydrogenolysis of the cyclopropyl group into an isopropyl group in the presence of a platinum catalyst and hydrobromic acid. *Tetrahedron*, *65*, 10623-10630, **2009**.
- (229) Hideaki Fujii, Ryo Ogawa, Kenji Ohata, Toru Nemoto, Mayumi Nagajima, Ko Hasebe, Hidenori Mochizuki, Hiroshi Nagase, Aerobic oxidation of indolomorphinan without the 4,5-epoxy bridge and subsequent rearrangement of the oxidation product to spiroindolinonyl-C-normorphinan derivative. *Bioorg. & Med. Chem.*, *17*, 5983-5988, **2009**.
- (228) Hideaki Fujii, Ryo Ogawa, Eikichi Jinbo, Saori Tsumura, Toru Nemoto, Hiroshi Nagase, Novel Oxidation Reaction of Tertiary Amines with Osmium Tetroxide. *Synlett*, *14*, 2341-2345, **2009**.
- (227) Yamamura S, Ohyama K, Nagase H, and Okada M, Zonisamide Enhances κ Receptor Associated Neurotransmitter Release in Striato-pallidal Pathway. *Neuropharmacology*, *57*, 322-331, **2009**.
- (226) Hideaki Fujii, Yoshikazu Watanabe, Yumiko Osa, Toru Nemoto, Noriko Sato, Hiroshi Nagase, Nobel Rearrangement Reaction of a 6,14-Endoethanomorphinan Derivative to a Benzomorphan Derivative. *Tetrahedron*, *65*, 4808-4813, **2009**.
- (225) Hiroshi Nagase, Toru Nemoto, Yumiko Osa, Hideaki Fujii, Masayuki Imai, Takashi Nakamura, Toshiyuki Kanemasa, Akira Kato, Hiroaki Gouda, Shuichi Hirono, Design and Synthesis of Novel δ Opioid Receptor Agonist and Their Pharmacologies. *Bioorg. & Med. Chem. Lett.*, *19*, 2792-2795, **2009**.
- (224) Akio Watanabe, Hideaki Fujii, Mayumi Nakajima, Ko Hasebe, Hidenori

- Mochizuki, Hiroshi Nagase, Synthesis of a new opioid ligand having the oxabicyclo[3.2.1]octane skeleton using a new rearrangement reaction. *Bioorg. & Med. Chem. Lett.*, *19*, 2416-2419, **2009**.
- (223) Hiroshi Nagase, Akio Watanabe, Masaya Harada, Mayumi Nakajima, Ko Hasebe, Hidenori Mochizuki, Kenji Yoza, Hideaki Fujii, Nobel Synthesis of a 1,3,5-Trioxazatriquinane Skeleton Using a Nitrogen Clamp. *Org. Lett.*, *11*, 539-542, **2009**.
- (222) Hideaki Fujii, Akio Watanabe, Toru Nemoto, Minoru Narita, Kan Miyoshi, Atsushi Nakamura, Tsutomu Suzuki, Hiroshi Ngase, Synthesis of novel twin drug consisting of 8-oxaendoethanotetrahydromorphides with a 1,4-dioxane spacer and its pharmacological activities: μ , κ , and ϵ opioid receptor antagonists. *Bioorg. & Med. Chem. Lett.*, *19*, 438-441, **2009**.
- (221) Toru Nemoto, Hideaki Fujii, Minoru Narita, Kan Miyoshi, Atsushi, Nakamura, Tsutomu Suzuki, Hiroshi Nagase, Synthesis of a novel 6, 14-epoxymorphinan derivative and its pharmacology. *Bioor. & Med. Chem.Lett.*, *18*, 6398-6401, **2008**.
- (220) Hiroshi Nagase, Naoshi Yamamoto, Toru Nemoto, Kenji Yoza, Kenshu Kamiya, Shuichi Hirono, Shinobu Momen, Naoki Izumimoto, Ko Hasebe, Hidenori Mochizuki, and Hideaki Fujii, Synthesis of a Stable Iminium Salt and Propellane Derivatives. *J. Org. Chem.*, *73*, 8093-8096, **2008**.
- (219) Koji Kawai, Jun Hayakawa, Toru Miyamoto, Yoshifumi Imamura, Shinichi Yamane, Hisanori Wakita, Hideaki Fujii, Kuniaki Kawamura, Hirotohi Matsuura, Naoki Izumimoto, Ryosuke Kobayashi, Takashi Endo, Hiroshi Nagase, Design, synthesis, and structure-activity relationship of novel opioid -agonists. *Bioorg. & Med.Chem.*, *16*, 9188-9201, **2008**.
- (218) Hideaki Fujii, Yumiko Osa, Marina Ishihara, Shinichi Hanamura, Toru Nemoto, Mayumi Nakajima, Ko Hasebe, Hidenori Mochizuki, Hiroshi Nagase, Synthesis of N-isobutylnoroxymoxyphone from naltrexone by a selctive cyclopropane ring opening reaction, *Bioorg. & Med.Chem.Lett.*, *18*, 4978-4981, **2008**.
- (217) Hideaki, Fujii, Satomi Imaide, Akio Watanabe, Toru Nemoto, Hiroshi Nagase, Novel Cleavage Reaction of the C16-N17 Bond in Naltrexone Derivatives. *Tetrahedoron Lett.*, *49*, 6293-6296, **2008**.
- (216) Satoshi Sakami, Masayuki Maeda, Koji Kawai, Takumi Aoki, Kuniaki Kawamura, Hideaki Fujii, Ko Hasebe, Mayumi Nakajima, Takashi Endo, Shinya Ueno, Tsuyoshi Ito, Junzo Kamei, and Hiroshi Nagase, Design and Synthesis of a Metabolically Stable and Potent Antitussive Agent, a Novel δ Opioid Receptor Antagonist, TRK-851. *Bioorg. & Med. Chem.*, *16*, 7956-7967,

2008.

- (215) Satoshi Sakami, Masayuki Maeda, Takumi Aoki, Kuniaki Kawamura, Koji Kawai, Hideaki Fujii, Takashi Endo, Shinya Ueno, Junzo Kamei, Ko Hasebe, Mayumi Nakajima, Tuyoshi Ito and Hiroshi Nagase, Structure-Antitussive Activity Relationships of Naltrindole Derivatives. Identification of Novel and Potent Antitussive Agents. *J. Med. Chem.* **51**, 4404-4411, **2008**.
- (214) Toru Nemoto, Hideaki Fujii, Minoru Narita, Kan Miyoshi, Atsushi Nakamura, Tsutomu Suzuki and Hiroshi Nagase, Synthesis of 4,6'-Epoxy-morphinan Derivatives and Their Pharmacologies. *Bioorg. & Med. Chem.*, **16**, 4304-4312, **2008**.
- (213) Toru Nemoto, Hideaki Fujii, Noriko Sato and Hiroshi Nagase, Synthesis of a novel 4,6'-epoxy-morphinan derivative and a highly strained novel conjugated ketone. *Tetrahedron Lett.*, **48**, 7413-7417, **2007**.
- (212) Yumiko Osa, Yoshihiro Ida, Hideaki Fujii, Toru Nemoto, Ko Hasebe, Shinobu Momen, Hidenori Mochizuki, and Hiroshi Nagase. Catalytic Aerobic Oxidation of nor-Binaltrephimine(nor-BNI) Analogs without 4,5-Epoxy Bridge Affords a More Selective Ligands for kappa Opioid Receptor than the Representative kappa Antagonist nor-BNI. *Chem.Pharm.Bull.* **55(10)** 1489-1493, **2007**.
- (211) Hiroshi Nagase, Akio Watanabe, Toru Nemoto, Naoshi Yamamoto, Yumiko Osa, Noriko Sato, Kenji Yoza and Toshitsugu Kai, Synthesis of opioid ligands having oxabicyclo[2.2.2]octane and oxabicyclo[2.2.1]heptane skeletons. *Tetrahedron Lett.*, **48**, 2547-2553, **2007**.
- (210) Y. Osa, Y. Ida, Y. Yano, K. Furuhashi, H. Nagase, A New Useful Conversion Method of Naltrexone to 14-Deoxynaltrexone. *Heterocycles.* **69**, 271-282, **2006**.
- (209) T. Mori, M. Nomura, K. Yoshizawa, H. Nagase, T. Sawaguchi, M. Narita, T. Suzuki, Generalization of NMDA-receptor antagonists to the discriminative stimulus effects of κ -opioid receptor agonists U-50,488H, but not TRK-820 in rats. *J. Pharmacol. Sci.*, **100(2)**, 157-161, **2006**.
- (208) Akio Watanabe, Toshitsugu Kai, and Hiroshi Nagase. Nobel Synthesis of the Ortho Ester Derivative of 4,5-Epoxy-morphinan. *Org. Lett.*, **8**, 523-526, **2006**.
- (207) Ariff Nurrochmad, Kishimoto Yayoi, Ozaki Masahiko, Yoshinori Yajima, Minoru Narita, Hiroshi Nagase, and Tsutomu Suzuki. Attenuation of the Morphine-Induced Rewarding Effect under Chronic Inflammatory Pain-like State in Rats. *Malaysian J. Pharmaceutical Sciences*, **3**, **1**, 11-24, **2005**.
- (206) NV. Solenkova, LN. Maslov, EV. Lishmanov Alu, P. Oeltgen, M. Govindaswami, ZhD. Bepalova, MV. Ovechinnikov, H. Nagase. Comparative study of the

- antiarrhythmic activity of μ - and δ -opioid receptor agonists during acute cardiac ischemia and reperfusion models in rats. *Eksp. Klin. Farmakol.*, 68(6), 25-29, **2005**.
- (205) N. Hirose, K. Murakawa, Y. Oi, T. Suzuki, H. Nagase, A. R. Cools and N. Koshimawa. Interactions among μ - and δ -Opioid Receptors, Especially Putative δ_1 - and δ_2 -Opioid Receptors, Promote Dopamine Release in the Nucleus accumbens. *Neuroscience*, 135, 213-225, **2005**.
- (204) Yumiko Osa, Yuka Hikima, Yuko Sato, Kouichi Takino, Yoshihiro Ida, Shuichi Hirono, and Hiroshi Nagase. Convenient Synthesis of Oxazolidinones by the Use of Halomethyloxirane, Primary Amine, and Carbonate Salt. *J Org. Chem.* 70, 5737- 5740, **2005**.
- (203) M. Narita, H. Akai, T. Kita, Y. Nagumo, M. Narita, N. Sunagawa, C. Hara, K. Hasebe, H. Nagase and T. Suzuki. Involvement of Mitogen-Stimulated P70-S6 Kinase in The Development of Stimulation to The Methamphetamine-Induced Rewarding Effect in Rats. *Neuroscience*, 132, 553-560, **2005**.
- (202) K. Fuse, I. Takahashi, S. Watanabe, Y. Aono, H. Ikeda, T. Saigusa, H. Nagase, T. Suzuki, N. Koshikawa and A. R. Cools. The Non-Peptidid δ opioid receptor Agonist TAN-67 Enhances Dopamine Efflux in The Nucleus Accumbens of Freely Moving Rats via a Mechanism that Involves both Glutamate and Free Radicals. *Neuroscience*, 130, 745-755, **2005**.
- (201) Hirokazu Mizoguchi, Leitermann, R. J., Minoru Narita, Hiroshi Nagase, Tsutomu Suzuki, Tseng, L. F. Region-dependent G-protein activation by kappa-opioid receptor agonists in the mouse brain. *Neurosci. Lett.*, 356, 145-147, **2004**.
- (200) Hasebe K, Kawai K, Suzuki T, Kawamura K, Tanaka T, Narita M, Nagase H, Suzuki T. Possible pharmacotherapy of the opioid κ receptor agonist for drug dependence. *Ann. NY Acad. Sci.*, 1025, 404-413, **2004**.
- (199) A. A. Platonov, T. V. Lasukova, L. N. Maslov, J. M. Downey, H. Nagase, M. V. Ovchinnikov. Opioid receptor agonists prevent the irreversible damage of Cardiomyocytes in ischemized reperfused rat heart. *Experimental & Clinical Pharmacology*, 67(6), 26-29, **2004**.
- (198) L. N. Maslov, T.V. Lasukova, Zh. D. Bepalova, P. Oeltgen, K. C. Rice, H. Nagase. δ -opioid receptor agonists are exhibited properties of partial agonists of δ -receptor in the experiment with isolated perfused heart. *Bulletin of Experimental Biology and Medicine* 138(10), 376-379, **2004**.
- (197) M Narita, H. Akai, Y. Nagumo, N. Sunagawa, K. Hasebe, H. Nagase, T. Kita, C.

- Haraand, T. Suzuki. Implications of Protein Kinase C in the Nucleus Accumbens in the Development of Sensitization to Methamphetamine in Rats. *Neuroscience*, *127*, 941-948, **2004**.
- (196) Yoshio Wakasa, Atsushi Fujiwara, Hideo Umeuchi, Takashi Endoh, Kiyoshi Okano, Toshiaki Tanaka, Hiroshi Nagase. Inhibitory effects of TRK-820 on systemic skin scratching induced by morphine in rhesus monkeys. *Life Sciences*, *24*, 2947-2957, **2004**.
- (195) Tomohisa Mori, Mutsuko Nomura, Kazumi Yoshizawa, Hiroshi Nagase, Minoru Narita, Tsutomu Suzuki. Differential properties between TRK-820 and U-50488H on the Discriminative stimulus effects in rats. *Life Sciences*, *75*, 2473-2482, **2004**.
- (194) Akiyoshi Saitoh, Yuji Kimura, Tomohiko Suzuki, Koji Kawai, Hiroshi Nagase, and Junzo Kamei. Potential Anxiolytic and Antidepressant-Like Activities of SNC80, a Selective δ -Opioid Agonist, in Behavioral Models in Rodents. *J. Pharmacol. Sci.*, *95*, 374-380, **2004**.
- (193) Hideaki Fujii, Minoru Narita, Hirokazu Mizoguchi, Junichi Hirokawa, Koji Kawai, Toshiaki Tanaka, Leon F. Tseng and Hiroshi Nagase. Rational drug design and synthesis of a selective ϵ -opioid receptor antagonist on the basis of the accessory site concept. *Bioorganic & Medicinal Chemistry Letters*, *14/16* 4241-4243, **2004**.
- (192) Hideaki Fujii, Minoru Narita, Hirokazu Mizoguchi, Miho Murachi, Toshiaki Tanaka, Koji Kawai, Leon F. Tseng and Hiroshi Nagase. Drug design and synthesis of ϵ opioid receptor agonist: 17-(cyclopropylmethyl)-4,5 α -epoxy-3,6, β -dihydroxy-6,14-endoethenomorphinan-(*N*-methyl-*N*-phenethyl)carboxamide(TAN-821) inducing antinociception mediated by putative ϵ -opioid receptor. *Bioorganic & Medicinal Chemistry*, *12*, 4133-4145, **2004**.
- (191) Hiromasa Horikiri, Noriyuki Hirano, Yoichiro Tanaka, Junji Oishi, Hitoshi Hatakeyama, Kuniaki Kawamura, and Hiroshi Nagase. Syntheses of 10-Oxo, 10 α -Hydroxy Derivatives of a Potent κ -Opioid Receptor Agonist, TRK-820. *Chem. Pharm. Bull.* *52(6)* 664-669, **2004**.
- (190) Kuniaki Kawamura, Hiromasa Horikiri, Jun Hayakawa, Chie Seki, Ken-ichi Yoshizawa, Hideo Umeuchi, and Hiroshi Nagase. Syntheses of Potential Metabolite of a Potent κ -Opioid Receptor Agonist, TRK-820. *Chem. Pharm. Bull.* *52(6)* 670-674, **2004**.
- (189) Hideaki Fujii, Noriyuki Hirano, Hiromi Uchiro, Kuniaki Kawamura, and

- Hiroshi Nagase. The First Example of the Stereoselective Synthesis of 7 α -Carbamoyl-4,5 α -epoxymorphinan via a Nobel and Reactive γ -Lactone. *Chem. Pharm. Bull.* 52(6) 747-750, **2004**.
- (188) Satoshi Matsuzaki, Hiroko Ikeda, Gaku Akiyama, Michiko Sato, Shoko Moribe Tsutomu Suzuki, Hiroshi Nagase, Alexander R. Cools, Noriaki Koshikawa. Role of μ - and δ -opioid receptors in the nucleus accumbens in turning Behaviour of rats. *Neuropharmacology* 46, 1089-1096, **2004**.
- (187) Kaoru Murakawa, Noriya Hirose, Koji Tanaka, Tsutomu Suzuki, Hiroshi Nagase Alexander R. Cools, Noriaki Koshikawa. Deltorphin II enhances extracellular levels of dopamine in the nucleus accumbens via opioid receptor-independent mechanisms. *Eur. J. pharmacol.*, 491, 31-36, **2004**.
- (186) Ichiro Takasaki, Tomohiko Suzuki, Atsushi Sasaki, Toshiaki Tanaka, Hiroshi Nagase, Kimiyasu Shiraki, Hiroshi Nijima, Yasushi Kuraishi. Suppression of acute herpetic pain-related responses by κ -opioid receptor agonist TRK-820 in mice. *J. Pharmacol. Exp. Ther.* 309(1), 36-41, **2004**.
- (185) Tomohiko Suzuki, Naoki Izumimoto, Yuko Takezawa, Morihiro Fujimura, Yuko Togashi, Hiroshi Nagase, Toshiaki Tanaka, Takashi Endo. Effect of repeated administration of TRK-820, a κ -opioid receptor agonist, on tolerance to its antinociceptive and sedative actions. *Brain Res.* 995, 167-175, **2004**.
- (184) Okura T, Varga EV, Hosohata Y, Navratilova E, Cowell SM, Rice K, Nagase H, Hruby VJ, Reske WR, Yamamura HI. Agonist-specific down regulation of the human δ -opioid receptor. *Eur. J. Pharmacol.* 459(1), 9-16, **2003**.
- (183) Hirokazu Mizoguchi, Hsiang-En Wu, Minoru Narita, Ichiro Sora, F. Scott Hall, George R. Uhl, Horace H. Loh, Hiroshi Nagase, and Leon F. Tseng. Lack of μ -Opioid Receptor-Mediated G-Protein Activation in the Spinal Cord of Mice Lacking Exon1 or Exon 2 and 3 of the Mor-1 Gene. *J. Pharmacol Sci* 93, 423-429, **2003**.
- (182) Junzo Kamei, Kayo Morita, Akiyoshi Saitoh, Hiroshi Nagase. The Antitussive Effects of Endomorphin-1 and Endomorphin-2 in Mice. *Eur. J. Pharmacol.*, 467, 219-222, **2003**.
- (181) Hsiang-En Wu, Moses Darpolor, Hiroshi Nagase, Leon F. Tseng. Acute antinociceptive tolerance and partial cross-tolerance to endomorphine-1 and endomorphine-2 given intrathecally in the mouse. *Neurosci. Lett.*, 348, 139-142, **2003**.
- (180) Hideo Umeuchi, Yuko Togashi, Toshiyuki Honda, Kaoru Nakao, Kiyoshi Okano, Tshiaki Tanaka, Hiroshi Nagase. Involvement of cenral μ -opioid system in the

- scratching behavior in mice, and the suppression of it by the activation of κ -opioid system. *Eur. J. Pharmacol.*, 477, 29-35, **2003**.
- (179) Masahiro Ohsawa, Hirokazu Mizoguchi, Minoru Narita, Hiroshi Nagase, Nae J. Dun, Lean F. Tseng. Involvement of β -arrestin-2 in modulation of the spinal antinociception induced by μ -opioid agonists in mouse, *Neurosci. Lett.*, 346, 13-16, **2003**.
- (178) Hirokazu Mizoguchi, Amanda Spaulding, Randy Leitermann, Hsiang-En Wu, Hiroshi Nagase, and Leon Tseng. Buprenorphine Blocks ϵ - and μ -Opioid Receptor-Mediated Antinociception in the Mouse, *J. Pharmacol. Ther. Exp.*, 306, 394-400, **2003**.
- (177) Hiroshi Tanaka, Hiroshi Ohno, Kuniaki Kawamura, Atsushi Ohtake, Hiroshi Nagase, and Takashi Takahashi. Solid-Phase Synthesis of Naltrindole Derivatives Using Fisher Indole Synthesis Based on One-Pot Release and Cyclization Methodology. *Org. Lett.*, 5, 8, 1159-1162, **2003**.
- (176) Hirokazu Mizoguchi, Kuei-Hung, Randy Leitermann, Minoru Narita, Hiroshi Nagase, Tsutomu Suzuki, Leon F. Tseng. Blockade of μ -Opioid Receptor-Mediated G-Protein Activation and Antinociception by TRK-820 in Mice, *Eur. J. Pharmacol.*, 461, 35-39, **2003**.
- (175) Hiroshi Ohno, Kuniaki Kawamura, Atsushi Otake, Hiroshi Nagase, Hiroshi Tanaka, Takashi Takahashi. Solid-Phase Synthesis of 6-Sulfonyl-amino Morphinan Libraries. *Synlett*, 93-96, **2002**.
- (174) Hirokazu Mizoguchi, Hsiang-En Wu, Minoru Narita, Horace H. Loh, Hiroshi Nagase, Leon F. Tseng. Loss of μ -opioid receptor-mediated G-protein activation in the pons/medulla of mice lacking the exons 2 and 3 of μ -opioid receptor gene. *Neurosci. Lett.*, 335, 91-94, **2002**.
- (173) H. Mizoguchi, H.E. WU, M. Narita, F.S. Hall, I. Sora, G. R. UHL, H. Nagase and L. F. Tseng. Antagonistic Property of Putative ϵ -Opioid Receptor-Mediated G-Protein Activation by beta-Endorphin in Pons/Medulla of the μ -Opioid Receptor Knockout Mouse. *Neuroscience*, 115, 715-721, **2002**.
- (172) Tomohisa Mori, Mutsuko Nomura, Hiroshi Nagase, Minoru Narita, Tsutomu Suzuki. Effects of a newly synthesized kappa-opioid receptor agonist, TRK-820, on the discriminative stimulus and rewarding effects of cocaine in rats. *Psychopharmacology* 161: 17-22, **2002**.
- (171) Yuya Ise, Minoru Narita, Hiroshi Nagase, Tsutomu Suzuki. Modulation of κ -opioidergic systems on mecamylamine-precipitated nicotine-withdrawal aversion in rats. *Neurosci. Lett.*, 323, 164-166, **2002**.

- (170) Hsiang-en Wu, Kuei-cheun Hung, Hirokazu Mizoguchi, Hiroshi Nagase, and Leon F. Tseng. Role of Endogenous Opioid Peptide in Modulation of Nocifensive Response to Formalin. *J. Pharmacol. Exp. Ther.* 300, 647-654, **2002**.
- (169) Yuko Togashi, Hideo Umeuchi, Kiyoshi Okano, Naoki Ando, Yoshitaka Yoshizawa, Toshiyuki Honda, Kuniaki Kawamura, Takashi Endoh, Jun Utsumi, Junzo Kamei, Toshiaki Tanaka, Hiroshi Nagase. Antipruritic Activity of the κ -Opioid Receptor Agonist, TRK-820. *Eur. J. Pharmacol.*, 435, 259-267, **2002**.
- (168) Ryan M. Fryer, Yigang Wang, Anna K. Hsu, Hiroshi Nagase, Garrwtt J. Gross. Dependence of delta1-opioid receptor-induced cardioprotection on a tyrosine kinase-dependent but not a Src-dependent pathway. *J. Pharmacol. Exp. Ther.*, 299, 477-482, **2001**.
- (167) Hosohata Y., Varga EV, Stropova D, Li X, Knapp RJ, Hrubby VJ, Rice KC, Nagase H, Roeske WR, Yamamura HI.. Mutation W284L of the Human δ opioid Receptor reveals agonist specific receptor conformations for G protein activation. *Life Sciences*, 68(19-20), 2233-2242, **2001**.
- (166) Masahiro Ohsawa, Maho Shiraki, Hirokazu Mizoguchi, Michiko Narita, Koji Kawai, Hiroshi Nagase, Eugene Y. Cheng, Minoru Narita, Leon F. Tseng. Release of [Met5]enkephalin from the spinal cord by intraventricularly administered endomorphin-2, but not endomorphin-1 in the anesthetized rat. *Neurosci. Lett.*, 316, 1-4, **2001**.
- (165) Minoru Narita, Satru Ozaki, Matsui Ioka, Hirokazu Mizoguchi, Hiroshi Nagase, Leon F. Tseng, Tsutomu Suzuki. Lack of the involvement of mu 1-opioid receptor subtype on motivational effects induced by the endogenous μ -opioid receptor ligands endomorphin-1 and -2 in the mouse. *Neurosci. Lett.* 308, 17-20, **2001**.
- (164) Hideaki Fujii, Koji Kawai, Kuniaki Kawamura, Akira Mizusuna, Yoshihiro Onoda, Miho Murachi, Toshiaki Tanaka, Takashi Endoh, and Hiroshi Nagase. Synthesis of Optically Active TAN-67, a Highly Selective δ - Opioid Receptor Agonist, and Investigation of its Pharmacological Properties. *Drug Design and Discovery*, 17, 325-330, **2001**.
- (163) Minoru Narita, Hirokazu Mizoguchi, Hiroshi Nagase, Tsutomu Suzuki, Leon F. Tseng. Up-Regulation of Spinal μ -Opioid Receptor Function to Activate G-Protein by Chronic Naloxone Treatment. *Brain Res.*, 913, 170-173, **2001**.
- (162) Junzo Kamei, Hiroshi Nagase. Norbinaltrphimine, a Selective κ -Opioid Receptor Antagonist, Induces an Itch-associated Response in Mice. *Eur. J. Pharmacol.*, 418, 141-145, **2001**.
- (161) Minoru Tsuji, Hiroshi Takeda, Teruhiko Matsumiya, Hiroshi Nagase, Minoru

- Narita, Tsutomu Suzuki. The Novel κ -Opioid Receptor Agonist TRK-820 Suppresses the Rewarding and Locomotor-Enhancing Effects of Morphine in Mice. *Life Sciences*, 68, 1717-1725, **2001**.
- (160) Minoru Narita, Hirokazu Mizoguchi, Hiroshi Nagase, Tsutomu Suzuki and Leon F. Tseng. Involvement of Spinal Protein Kinase C γ in the Attenuation of Opioid μ -Receptor-Mediated G-Protein Activation after Chronic Intrathecal Administration of [D-Ala²,N-MePhe⁴,Gly⁰-O⁵] Enkephalin. *J. Neurosci.* 21, 3715-3720, **2001**.
- (159) Masahiro Ohsawa, Hirokazu Mizoguchi, Minoru Narita, Hiroshi Nagase, John P.Kampaine, and Leon F. Tseng. Differential Antinociception Induced by Sponally Administered Endomorphin-1 and Endomorphin-2 in the Mouse. *J. Pharmacol. Exp. Ther.* 298, 592-597, **2001**.
- (158) Hiroshi Nagase, Yoshinori Yajima, Hideaki Fujii, Kuniaki Kawamura Minoru Narita, Junzo Kamei, Tsutomu Suzuki. The Pharmacological Profile of delta Opioid Receptor Ligands, (+) and (-) TAN-67 on Pain Modulation. *Life Sciences*. 68, 2227-2231, **2001**.
- (157) Minoru, Narita, Hirokazu Mizoguchi, Tomohiko Suzuki, Michiko Narita, Nae J. Dun, Satoshi Imai, Yoshinori Yajima, Hiroshi Nagase, Tsutomu Suzuki, and Leon F. Tseng. Enhanced μ -Opioid Responses in the Spinal Cord of Mice Lacking Protein Kinase C gamma Isoform. *J. Biological Chem.*, 276, 15409-15414, **2001**.
- (156) M. Narita, S. Ozaki, M. Ioka, H. Mizoguchi, H. Nagase, L.F. Tseng and T. Suzuki. Different Motivational Effects Induced by the Endogenous μ -Opioid Receptor Ligands Endomorphin-1 and -2 in the Mouse. *Neuroscience*, 105, 213-218, **2001**.
- (155) Takashi Endoh, Atsushi Tajima, Naoki Izumimoto, Tomohiro Suzuki, Akiyoshi Saitoh, Tsutomu Suzuki, Minoru Narita, Junzo Kamei, Leon F. Tseng, Hirokazu Mizoguchi and Hiroshi Nagase. TRK-820, a Selective κ -Opioid Agonist, Produces Potent Antinociception in Cynomolgus Monkeys. *Jpn. J. Pharmacol.* 85, 282-290, **2001**.
- (154) Joon Huh, Garratt J. Gross, Hiroshi Nagase, and Bruce T. Liang. Protection of cardiac myocytes via delta1-opioid receptors, protein kinase C, and mitochondrial KATP channels. *Am J. Physiol. Heart Circ. Physiol.*, 280:H377-383, **2001**.
- (153) Junzo Kamei, Ko Zushida, Hiroshi Nagase. Role of Cholecystokinin in the Reduction of Endomorphin-2-induced Antinociception in Diabetic Mice. *Eur. J. Pharmacol.*, 416, 95-99, **2001**.
- (152) Junzo Kamei, Ko Zushida, Masahiro Ohsawa, Hiroshi Nagase. The

- Antinociceptive Effects of Endomorphin-1 and Endomorphin-2 in Diabetic Mice. *Eur. J. Pharmacol.* 391, 91-96, 2000.
- (151) Yuya Ise, Minoru Narita, Hiroshi Nagase, Tsutomu Suzuki. Modulation of opioidergic System on Mecaylamine-Precipitated Nicotine-Withdrawal Aversion in Rats. *Psychopharmacology*, 151: 49-54, 2000.
- (150) Tsutomu Suzuki, Takeshi Aoki, Orié Ohnishi, Hiroshi Nagase, Minoru Narita. Different Effects of NMDA/group I Metabotropic Glutamate Receptor Agents in δ -, and μ -Opioid Receptor Agonist-Induced Supraspinal antinociception. *Eur. J. Pharmacol.*, 396, 23-28, 2000.
- (149) H. Mizoguchi, M. Narita, H.-E. WU, M. Narita, T. Suzuki, H. Nagase, and L.F. Tseng. Differential Involvement of μ 1-Opioid Receptors in Endomorphine- and β -Endorphin-Induced G-Protein Activation in the Mouse Pons/Medulla. *Neuroscience*, 100, 835-839, 2000.
- (148) Hirokazu Mizoguchi, Minoru Narita, Hiroshi Nagase, Leon F. Tseng. Activation of G-proteins in the mouse pons/medulla by beta-endorphin is mediated by the stimulation of μ - and putative epsilon-receptors. *Life Sci.*, 67, 2733-2743, 2000.
- (147) Hisanori Wakita, Hideo Yoshiwara, Yukishige Kitano, Hisao Nishiyama, and Hiroshi Nagase. Preparative Resolution of 2-Methyl-4-Hexynoic Acid for the Synthesis of Optically Active m-Phenylene PGI₂ Derivatives and Determination of their Absolute Configuration. *Tetrahedron Asymmetry*, 11, 2981-2989, 2000.
- (146) Masahiro Ohsawa, Hirokazu Mizoguchi, Minoru Narita, Mei Chu, Hiroshi Nagase, and Leon F. Tseng. Differential Mechanisms Mediating Descending Pain Controls for Antinociception Induced by Supraspinally Administered Endomorphin-1 and Endomorphin-2 in the Mouse. *J. Pharmacol. and Exp. Ther.*, 294, 1106-1111, 2000.
- (145) Galen M. Pieper, Hirokazu Mizoguchi, Masahiro Ohsawa, Junzo Kamei, Hiroshi Nagase, and Leon F. Tseng. Decreased Opioid-Induced Antinociception but Unaltered G-protein Activation in the Genetic-Diabetic NOD mouse. *Eur. J. Pharmacol.*, 401, 375-379, 2000.
- (144) Masahiro Ohsawa, Hirokazu Mizoguchi, Minoru Narita, Michiko Narita, Junzo Kamei, Hiroshi Nagase, Leon F. Tseng. Effects of a μ -Opioid Receptor Agonist on G-protein Activation in Streptozocin-Induced Diabetic Mice. *Eur. J. Pharmacol.*, 401, 55-58, 2000.
- (143) Minoru Tsuji, Mitsuaki Yamazaki, Hiroshi Takeda, Teruhiko Matsumiya, Hiroshi Nagase, Leon F. Tseng, Minoru Narita, Tsutomu Suzuki. The Novel κ -Opioid Receptor Agonist TRK-820 has no effect on the development of

- tinociceptive Tolerance to Morphine in Mice. *Eur. J. Pharmacol.*, 394, 91-95, 2000.
- (142) Minoru Tsuji, Hiroshi Takeda, Teruhiko Matsumiya, Hiroshi Nagase, Mitsuaki Yamazaki, Minoru Narita and Tsutomu Suzuki. A Novel kappa-Opioid Receptor Agonist, TRK-820, Block the Development of Physical Dependence on Morphine in Mice. *Life Sci.*, 66, 353-358, 2000.
- (141) Ryan Fryer, Anna K. Hsu, Hiroshi Nagase, Garrett J. Gross. Opioid-Induced Cardioprotection Against Myocardial Infarction and Arrhythmias: Mitochondrial versus Sarcolemmal KATP Channels, *J. Pharmacol. Exp. Ther.* 294, 451-45, 2000.
- (140) Yoshinori Yajima, Minoru Narita, Makoto Tsuda, Satoshi, Imai Junzo Kamei Hiroshi Nagase and Tsutomu Suzuki. Modulation of NMDA- and (+)TAN-67-Induced Nociception by GABAB Receptor the Mouse Spinal Cord. *Life Sciences*, 68, 719-725, 2000.
- (139) Hisanori Wakita, ideo Yoshiwara, isao Nishiyama and Hiroshi Nagase. Total Synthesis of Optically Active m-Phenylene PGI₂ Derivative. Beraprost. *Heterocycle* 53, 1085-1110, 2000.
- (138) Takashi Endoh, Atsushi Tajima, Tomohiko Suzuki, Junzo Kamei, Tsutomu Suzuki, Minoru Narita, Leon Tseng and Hiroshi Nagase. Characterization of the Antinociceptive Effects of TRK-820 in the Rat. *Eur J. Pharmacol.* 387, 133-140, 2000.
- (137) Yoshinori Yajima, Minoru Narita, Yuki Takahashi-Nakano, Miwa Misawa, Hiroshi Nagase, Hirokazu Mizoguchi, Leon F. Tseng and Tsutomu Suzuki. Effect of Differential Moduration of μ , δ , and κ -Opioid System on the Bicuculline-Induced Convulsion in the Mouse. *Brain Research*, 862, 120-126, 2000.
- (136) Minoru Narita, Hirokazu Mizoguchi, Michiko Narita, Nae J. Dun, Bang H. Hwang, Takashi Endoh, Tomohiko Suzuki, Hiroshi Nagase, Tsutomu Suzuki. John P. Kampine and Leon F. Tseng. G-protein Activation by Endomorphins in the mouse Periaqueductal Gray Matter. *J. Biomed. Science*, 7, 221-225, 2000.
- (135) Leon F. Tseng, Minoru Narita, Chiaki Suganuma, Hirokazu Mizoguchi, Hiroshi Nagase and John P. Kampine. Differential Antinociceptive Effects of Endomorphin-1 and Endomorphin-2 in the Mouse. *J. Pharmacol. Exp. Ther.* 292, 576-583, 2000.
- (134) Junzo Kamei, Masahiro Ohsawa, Takako Kashiwazaki, Hiroshi Nagase. Antinociceptive effects of the ORL1 receptor agonist nociceptin/orphanin FQ in diabetic mice. *Eur. J. Pharmacol.*, 370, 109-116, 1999.

- (133) Junzo Kamei, Takako Kashiwazaki, Kentaro Taki, Hideki Hitosugi, Hiroshi Nagase. Allogenic mediator-induced nociceptive response in diabetic mice. *Eur. J. Pharmacol.* 369, 319-323, **1999**.
- (132) Junzo Kamei, Kentaro Taki, Masahiro Ohsawa, Hiroshi Nagase, Kenji Onodera. Involvement of spinal delta 1-opioid receptors in forced walking stress-induced antinociception in the tail-flick test in mice. *Methods Find. Exp. Clin. Pharmacol.*, 21, 595-598, **1999**.
- (131) Junzo Kamei, Masahiro Ohsawa, Tomohiko Suzuki, Akiyoshi Saitoh, Takashi Endoh, Minoru Narita, Leon F. Tseng, Hiroshi Nagase. The Modulatory Effect of (+)-TAN-67 on the antinociceptive Effects of the Nociceptin/Orphan FQ in Mice. *Eur. J. Pharmacol.* 383, 241-247, **1999**.
- (130) Takashi Endoh, Hirotohi Matsuura, Atsushi Tajima, Naoki Izumimoto, Chiko Tajima, Tomohiko Suzuki, Akiyoshi Saitoh, Tsutomu Suzuki, Minoru Narita, Leon Tseng and Hiroshi Nagase. Potent Antinociceptive Effects of TRK-820, A Novel κ -Opioid Receptor Agonist. *Life Sci.*, 65, 1685-1694, **1999**.
- (129) Hisanori Wakita, Hideo Yoshiwara, Atsuko Tajima, Yukishige Kitano, and Hiroshi Nagase. Preparative Resolution of a Key Intermediate for the Synthesis of Optically Active *m*-Phenylene PGI₂ Derivative. *Tetrahedron Asymmetry.*, 10, 4099-4105, **1999**.
- (128) Minoru Narita, Hirokazu Mizoguchi, David E. Oji, Michiko Narita, Nae J. Dun, Bang H. Hwang, Hiroshi Nagase and Leon F. Tseng. Identification of the Opioid like Orphan G-Protein-Coupled Receptor System in the Mouse Spinal Cord. *British J. Pharmacol.* 128, 1300-1306, **1999**.
- (127) Takahiro Seki, Shinichiro Awamura, Chiaki Kimura, Soichiro Ide, Kyoko Sakano, Masabumi Minami, Hiroshi Nagase, Masamichi Satoh. Pharmacological Properties of TRK-820 on Cloned μ , δ , κ -Opioid Receptors and Nociceptin Receptor. *Eur. J. Pharmacol.* 376, 159-167, **1999**.
- (126) Ryan M. Fryer, Anna K. Hsu, Janis T. Eells, Hiroshi Nagase, Garrett J. Gross. Opioid-Induced Second Window of Cardioprotection Potential Role of Mitochondrial KATP Channels. *Circulation Res.*, 84, 846-851, **1999**.
- (125) Minoru Tsuji, Minoru Narita, Hirokazu Mizoguchi, Michiko Narita, Masahiro Ohsawa, Junzo Kamei, Hiroshi Nagase, Hiroshi Takeda, Teruhiko Matsumiya and Leon F. Tseng. Region-Dependent G-Protein Activation by μ -, δ_1 - and δ_2 -Opioid Receptor Agonists in the Brain: Comparison between the Midbrain and Forebrain. *Life Sci.* 65, PL 233-239, **1999**.
- (124) H. Mizoguchi, M.Narita, D.E.Oji, C. Suganuma, H. Nagase, I. Sora, G.R.UHL,

- E.Y.Cheng, and L. F.Tseng. The μ -opioid Receptor Gene-Dose/Dependent Reduction in G-Protein Activation in the Pons/Medulla and Antinociception induced by endomorphins in μ -Opioid Receptor Knockout Mice. *Neuroscience*, *94*, 203-207, **1999**.
- (123) S. Matsuzawa, T. Suzuki, M. Misawa and H. Nagase. Involvement of dopamine D₁ and D₂ receptors in the ethanol-associated place preference in rats exposed to conditioned fear stress. *Brain Res.* *835*, 298-305, **1999**.
- (122) S. Matsuzawa, T. Suzuki, M. Misawa and H. Nagase. Role of 5-HT₃ and opioid receptors in the ethanol-induced place preference in rats exposed to conditioned fear stress. *Life Sci.*, *64*, (21), PL241-249, **1999**.
- (121) S. Matsuzawa, T. Suzuki, M. Misawa and H. Nagase. Different roles of μ -, δ - and κ -opioid receptors in the ethanol-associated place preference in rats exposed to conditioned fear stress. *Eur. J. Pharmacol.*, *368*, 9-16, **1999**.
- (120) Ping Xiao, Hajime Kubo, Masahiro Ohsawa, Kimio Higashiyama, Hiroshi Nagase, Yu-Ning Yan, Jia-Shi Li, Junzo Kamei, and Shigeru Ohomiya. κ -Opioid Receptor-Mediated Antinociceptive Effects of Stereoisomers and Derivatives of (+)-Materine in Mice. *Planta Medica*, *65*, 230-233, **1999**.
- (119) Hisanori Wakita, Kazuhisa Matsumoto, Hideo Yoshiwara, Hisao Nishiyama, and Hiroshi Nagase. Synthesis of 5,6,7 -Trinor-4,8-Inter-m-Phenylene PGI₂ and Beraprost. *Tetrahedron*, *55*, 2449-2474, **1999**.
- (118) T. Suzuki, Y. Kishimoto, M. Misawa, H. Nagase and F. Takeda. Role of the κ -opioid system in the attenuation of the morphine-induced place preference under chronic pain. *Life Sci.* *64*(1), PL1-7, **1998**.
- (117) Masahiro Ohsawa, Hiroshi Nagase and Junzo Kamei. Role of Intracellular Calcium in Modification of Mu and Delta Opioid Receptor-Mediated Antinociception by Diabetes in Mice. *J. Pharmacol. Exp. Ther.*, *286*, 780-787, **1998**.
- (116) S. Matsuzawa, T. Suzuki, M. Misawa and H. Nagase. Involvement of μ - and δ -opioid receptors in the ethanol-associated place preference in rats exposed to foot shock stress. *Brain Res.* *803*(1/2), 169-177, **1998**.
- (115) Hisanori Wakita, Yutaka Hosono, and Hiroshi Nagase. Regioselective Protection of 11-Hydroxy Group and Inversion of Configuration at C-15 in m-Phenylene PGI₂ Derivatives. *Heterocycles*, *48*, 2559-2571, **1998**.
- (114) Hiroshi Nagase, Koji Kawai, Jun Hayakawa, Hisanori Wakita, Akira Mizusuna, Hirotohi Matsuura, Chiko Tajima, Yuko Takezawa, and Takashi Endoh. Rational Drug Design and Synthesis of a Highly Selective Nonpeptide δ -Opioid

- Agonist,(4aS*,12aR*)-4a-(3-Hydroxyphenyl)-2-methyl-1,2,3,4,4a,5,12,12a-Octahydro-1,2,3,4-benzopyrido[3,4-b]acridine (TAN-67). *Chem. Pharm. Bull.* **46**, (11) 1695-1702, **1998**.
- (113) Hisao Nishiyama, Naoya Sakata, Hayato Motoyama, Hisanori Wakita, and Hiroshi Nagase. Synthesis of Optically Active Cyclopenta[b]benzofuran as a Precursor of m-Phenylene-prostacyclin. *Synlett*, 930-932, **1998**.
- (112) Minoru Narita, Hirotochi Mizoguchi, George S. Oji, Esther L. Tseng, Chiaki Suganuma, Hiroshi Nagase and Leon F. Tseng. Characterization of Endomorphin-1 and -2 on [35S] GTP γ S Binding in the mouse Spinal Cord. *Eur. J. Pharmacol.*, **351**,383-387, **1998**.
- (111) T.H.Burkey, F.J.Ehlert, Y. Hosohata, R.M.Quock, S.Cowell, K. Hosohata, E.Varga, D. Stroova, X.Li,C.Slate, H.Nagase, F.Porraeca, V.J.Hruby, W.R.Roeske and H.I.Yamamura. The Efficacy of κ -Opioid Receptor-Selective Drugs. *Life Sciences*, **62**, 1531-1536, **1998**.
- (110) Koji Kawai, Kuniaki Kawamura and Hiroshi Nagase. Establishment of Efficient Hydroxylation at Angular Position of 4a-Aryl-trans-Deca-Hydroisoquinolines. *Heterocycles*, **48**, No 5, 949-959, **1998**.
- (109) Jo El. Schultz, Anna K. Hsu, Hiroshi Nagase, and Garrrwt J. Gross. TAN-67, A Delta1-Opioid Receptor Agonist, Reduces Infarct Size via Activation of Gi/o Proteins and KATP Channels. *Am. Physiol. Soc.*, H909-H914, **1998**.
- (108) Hiroshi Nagase, Jun Hayakawa, Kuniaki Kawamura, Koji Kawai, Yuko Takezawa, Hirotochi Matsuura, Chiko Tajima, and Takashi Endoh. Discovery of a Structurally Novel Opioid κ -Agonist Derived from 4,5-Epoxy Morphinan. *Chem Pharm. Bull.* **46(2)**, 366-369, **1998**.
- (107) Kuniaki Kawamura, Koji Kawai, Toru Miyamoto, Koji Ooshima, and Hiroshi Nagase. Synthesis of trans-4a-Aryl-6-Oxodekahydro-isoquinolines. *Heterocycle*, **48**, 267-283, **1998**.
- (106) Junzo Kamei, Takako Kashiwazaki, Hideki Hitosugi, Hiroshi Nagase. The role of spinal delta1-opioid receptors in inhibiting the formalin-induced nociceptive response in diabetic mice. *Eur. J. Pharmacol.*, **326**, 31-36, **1997**.
- (105) Junzo Kamei, Midori Sodeyama, Makoto Tsuda, Tsutomu Suzuki, Hiroshi Nagase. Antinociceptive effect of buprenorphine in mu1-opioid receptor deficient CXBK mice. *Life Sci.*, **60**, PL 333-337, **1997**.
- (104) Minoru Narita, Hirokazu Mizoguchi, Hiroshi Nagase, Leon F. Tseng. Use of Antisense Oligodeoxynucleotide to δ -Opioid Receptor m-RNA in the Study of δ -Opioid Receptors in the Spinal Cord of the Mouse. *Psychopharmacology* **133**,

- 347-350, **1997**.
- (103) Hisao Nishiyama, Naoya Sakata, Yukihiro Motoyama, Hisanori Wakita, and Hiroshi Nagase. Asymmetric Induction in Intramolecular SN₂ Reaction. Enantioselective Synthesis of Cyclopenta[b]benzofuran with Chiral Lithium Alkoxides. *Synlett*, *10*, 1147-1148, **1997**.
- (102) Tsutomu Suzuki, Tomohisa Mori, Minoru Tsuji, Miwa Misawa, and Hiroshi Nagase. The Role of δ -Opioid Receptors in the Discriminative Stimulus Properties of a Low Dose of Methamphetamine. *Eur. J. Pharmacol.*, *331*, 1-8, **1997**.
- (101) Junzo Kamei, Ping Xiao, Masahiro Ohsawa, Hajime Kubo, Kimio Higashiyama, Hiroshi Takahashi, Jiashi Li, Hiroshi Nagase, and Shigeru Ohmiya. Antinociceptive Effects of (+)-Matrine in Mice. *Eur. J. Pharmacol.*, *337*, 223-226, **1997**.
- (100) Junzo Kamei, Masahiro Ohsawa, Tsutomu Suzuki, and Hiroshi Nagase. Modification of Morphine-Induced Place Preference by Diabetes. *Eur. J. Pharmacol.*, *337*, 137-145, **1997**.
- (99) Hideaki Fujii, Akira Mizusuna, Ryuji Tanimura, and Hiroshi Nagase. A Novel Abnormal Rearrangement in the Fisher Indole Synthesis. *Heterocycles*, *45*, No. 11, 2109-2112, **1997**.
- (98) Tsutomu Suzuki, Minoru Tsuji, Tomohisa Mori, Miwa Misawa and Hiroshi Nagase. Involvement of δ_1 and δ_2 Opioid Receptor Subtypes in the Dependence on Morphine in Mice. *Pharmacol., Biochem. Behavior*, *57*, 293-299, **1997**.
- (97) Tsutomu Suzuki, Yayoi Kishimoto, Miwa Misawa and Hiroshi Nagase. The mechanism of suppression of morphine-induced rewarding effect under chronic pain. *Chronic Pain*, *16*, 121-125, **1997**.
- (96) Tsutomu Suzuki, Tomohisa Mori, Minoru Tsuji, Jun Maeda, Yayoi Kishimoto, Miwa Misawa, and Hiroshi Nagase. Differential effects of μ , δ and κ -Opioid Receptor Agonists on the Discriminative Stimulus Properties of Cocaine in Rats. *Eur. J. Pharmacol.*, *324*, 21-29, **1997**.
- (95) Tsutomu Suzuki, Minoru Tsuji, Tomohisa Mori, Hiroki Ikeda, Miwa Misawa, Hiroshi Nagase. Involvement of Dopamine-Dependent Mechanism in the Rewarding Effects Mediated by δ -Opioid Receptor Subtypes in mice. *Brain Res.*, *744*, 327-334, **1997**.
- (94) Raymond M. Quock, Yoshiaki Hosohata, Richard J. Knapp, Thomas H. Burkey, Keiko Hosohata, Xiaoyan Zhang, Kenner C. Rice, Hiroshi Nagase, Victor J. Hruby, Frank Porreca, William R. Roeske and Henry I. Yamamura. Relative

- Efficacies of δ -Opioid Receptor Agonists at the Cloned Human δ -Opioid Receptor. *Eur. J. Pharmacol.*, 326, 101-104, **1997**.
- (93) Leon F. Teng, Minoru Narita, Hirokazu Mizoguchi, Kohji Kawai, Akira Mizusuna, Junzo Kamei, Tsutomu Suzuki and Hiroshi Nagase. δ_1 Opioid Receptor-Mediated Antinociceptive Properties of Nonpeptidic Delta Opioid Receptor Agonist, (-)-TAN-67, in the Mouse Spinal cord. *J. Pharmacol. Exp. Ther.* 280, 600-605, **1997**.
- (92) Junzo Kamei, Koji Kawai, Akira Mizusuna, Akiyoshi Saitoh, Kayo Morita, Minoru Narita, Leon F. Tseng and Hiroshi Nagase. Spinal δ_1 -Opioid Receptor-Mediated Antinociceptive Properties of (-)-TAN-67 in Diabetic Mice. *Eur. J. Pharmacol.*, 322, 27-30, **1997**.
- (91) Junzo Kamei, Misahiro Ohsawa, Hiroshi Nagase. Possible involvement of μ_2 -opioid receptor-mediated mechanisms in morphine-induced enhancement of the pentobarbital-induced loss of the righting reflex in the mouse. *Life Sci.*, 59, PL349-353, **1996**.
- (90) Eva V. Varga, Xiaoping Li, Dagmar Strpova, Teresa Zalewska, Robert S. Landsman, Richard J. Nkapp, Ewa Malatynska, Kohji Kawai, Akira Mizusuna, Hiroshi Nagase, Silvia N. Calderon, Kenner Rice, Victor J. Hruby, William R. Roeske, and Henry Yamamura. The Third Extracellular Loop of the Human delta-Opioid Receptor Determines the Selectivity of δ -Opioid Agonists. *Molecular Pharmacol.*, 50, 1619-1624, **1996**.
- (89) Junzo Kamei, Tsutomu Suzuki, Hiroshi Nagase. Agonist and Antagonist Properties of Dihidromorphine for μ -Opioid Receptors in Mice. *Neuroscience Lett.*, 215, 87-90, **1996**.
- (88) Tsutomu Suzuki, Minoru Tsuji, Tomohisa Mori, Miwa Misawa, Takashi Endo and Hiroshi Nagase. Effect of Highly Selective and nonpeptide δ Opioid Receptor Agonist TAN-67 on the Morphine-Induced Place Preference in Mice. *J. Pharmacol., Exp., Ther.*, 279, 177-185, **1996**.
- (87) Junzo Kamei, Kayo Morita, Akiyoshi Saitoh, Tsutomu Suzuki and Hiroshi Nagase. Antinociceptive and Antitussive Effects of Morphine in the DA-Bg(Beige) Rat. *European J. Pharmacol.*, 300, 75-78, **1996**.
- (86) Makoto Tsuda, Tsutomu Suzuki, Miwa Misawa and Hiroshi Nagase. Involvement of the Opioid System in the Anxiolytic Effect of Diazepam in Mice. *Eur. J. Pharmacol.*, 307, 7-14, **1996**.
- (85) Tsutomu Suzuki, Minoru Tsuji, Tomohisa Mori, Miwa Misawa and Hiroshi Nagase. The Effects of Dopamine D1 and D2 Receptor Antagonists on the

- Rewarding Effects of δ_1 and δ_2 Opioid Receptor Agonists in Mice. *Psychopharmacology*, 124, 211-218, **1996**.
- (84) Hirokazu Mizoguchi, Minoru Narita, Hiroshi Nagase, Tsutomu Suzuki, Raymond M. Quock and Leon F. Tseng. Use of Antisense Oligonucleotide to Determine δ -Opioid Receptor Involvement [D-Ala] Deltorphine II-Induced Locomotor Hyperactivity, *Life Sci.*, 59, 69-73, **1996**.
- (83) Junzo Kamei, Tsutomu Suzuki, Miwa Misawa, Hiroshi Nagase, Yutaka Kasuya. Antinociceptive Effect of Dihydroetorphine in Diabetic Mice. *European J. Pharmacol.*, 275, 109-113, **1995**.
- (82) Junzo Kamei, Akiyoshi Saitoh, Tsutomu Suzuki, Miwa Misawa, Hiroshi Nagase and Yutaka Kasuya. Buprenorphine Exerts Aninociceptive Action via μ_1 -Opioid Receptors. *Life Sciences*, 56, 285-290, **1995**.
- (81) Junzo Kamei, Akiyoshi Saitoh, Kayo Morita, and Hiroshi Nagase. Antagonistic Effect of Buprenorphine on the Antitussive Effect of Morphine is Mediated via the Activation of μ_1 -Opioid Receptors. *Life Sciences*, 57, 231-235, **1995**.
- (80) Junzo Kamei, Masahiro Ohsawa, Akiyoshi Saitoh, Yuriko Iwamoto, Tsutomu Suzuki, Miwa Misawa, Hiroshi Nagase and Yutaka Kasuya. Modification of μ -Opioid Agonist-Induced Locomotor Activity and Development of Morphine Dependence by Diabetes. *J. Pharmacol. Exp. Therapeutics*, 274, 700-706, **1995**.
- (79) Tsutomu Suzuki, Minoru Tsuji, Tomohisa Mori, Miwa Misawa and Hiroshi Nagase. Effect of Naltrindole on the Development of Physical Dependence on Morphine in Mice: A Behavioral and Biochemical Study. *Life Sciences*, 57, 247-252, **1995**.
- (78) Tsutomu Suzuki, Tomohisa Mori, Minoru Tsuji, Miwa Misawa, Hiroshi Nagase. Discriminative Stimulus Properties of Morphine Mediated by μ_1 -Opioid Receptors. *European J. Pharmacol.*, 284, 195-198, **1995**.
- (77) Hirokazu Mizoguchi, Minoru Narita, Hiroshi Nagase, and Leon F. Tseng. Antisense Oligodeoxynucleotide to a δ -Opioid Receptor Blocks the Antinociception Induced by Cold Water Swimming. *Regulatory Peptide*, 59, 255-259, **1995**.
- (76) Richard J. Knapp, Robert Landsman, Sue Waite, Ewa Malatynska, Eva Varga, Wajujahal Haqt, Victor J. Hruby, William R. Roeske, Hiroshi Nagase, and Henry I. Yamamura. Properties of TAN-67, a Nonpeptidic δ -Opioid Agonist, at Cloned Human δ - and μ -Opioid Receptors. *European J. Pharmacol.*, 291 129-134, **1995**.
- (75) R. J. Knapp, R. Landsman, G. Santro, I.A. Deleon, E. Malatynska, E. Varga, H. Nagase, S.N. Calderon, K. Rice, F. Porecca, W. R. Roeske, and Henry I.

- Yamamura. Efficacy of Peptide and Nonpeptidic Agonists at the Cloned Human δ -Opioid Receptor. *Proc. West. Pharmacol. Soc.* 38, 141-143, **1995**.
- (74) Tsutomu Suzuki, Minoru Tsuji, Tomohisa Mori, Miwa Misawa, Takashi Endho, Hiroshi Nagase. Effects of a Highly Selective Nonpeptide δ -Opioid Receptor Agonist, TAN-67, on Morphine-Induced Antinociception in Mice. *Life Sciences*, 57, 155-168, **1995**.
- (73) Junzo Kamei, Akiyoshi Saitoh, Masahiro Ohsawa, Tsutomu Suzuki, Miwa Misawa, Hiroshi Nagase, and Yutaka Kasuya. Antinociceptive Effects of the Selective non-Peptidic δ -Receptor agonist TAN-67 in Diabetic Mice. *European J. Pharmacol.*, 276, 131-135, **1995**.
- (72) Junzo Kamei, Yuriko Iwamoto, Tsutomu Suzuki, Miwa Misawa, Hiroshi Nagase, Yutaka Kasuya. Antitussive effect of dihydroetorphine in mice. *Eur. J. Pharmacol.*, 260, 257-259, **1994**.
- (71) Junzo Kamei, Akiyoshi Saitoh, Yuriko Iwamoto, Masahiko Funada, Tsutomu Suzuki, Miwa Misawa, Hiroshi Nagase, Yutaka Kasuya. Effects of diabetes on spontaneous locomotor activity in mice. *Neurosci. Lett.*, 178, 69-72, **1994**.
- (70) Junzo Kamei, Masahiro Ohsawa, Miwa Misawa, Hiroshi Nagase, Yutaka Kasuya. Effects of diabetes on the morphine-induced Straub tail reaction in mice. *Neurosci. Lett.*, 178, 66-68, **1994**.
- (69) Junzo Kamei, Yuriko Iwamoto, Miwa Misawa, Hiroshi Nagase, Yutaka Kasuya. Involvement of adenosine A1 receptors in antitussive effect in mice. *Life Sci.*, 55, PL383-388, **1994**.
- (68) Tsutomu Suzuki, Michiharu Yoshiike, Hirokazu Mizoguchi, Junzo Kamei, Miwa Misawa, Hiroshi Nagase. Blockade of delta-opioid receptors prevents morphine-induced place preference in mice. *Jpn. J. Pharmacol.*, 66, 131-137, **1994**.
- (67) Tsutomu Suzuki, Tomohisa Mori, Minoru Tsuji, Miwa Misawa, Hiroshi Nagase. The role of delta-opioid receptor subtypes in cocaine- and methamphetamine-induced place preferences. *Life Sci.*, 55, PL339-344, **1994**.
- (66) Junzo Kamei, Yuriko Iwamoto, Hideki Hitosugi, Miwa Misawa, Hiroshi Nagase, Yutaka Kasuya. Differential mediation of cold water swim stress-induced antinociception by delta-opioid receptor subtypes in diabetic mice. *Life Sci.*, 54, PL425-430, **1994**.
- (65) Junzo Kamei, Yuriko Iwamoto, Miwa Misawa, Hiroshi Nagase, Yutaka Kasuya. Evidence for differential modulation of mu-opioid receptor-mediated antinociceptive and antitussive activities by spleen-derived factor(s) from

- diabetic mice. *Neuropharmacology*, 33, 1553-1558, **1994**.
- (64) Tstomu Suzuki, Tomohisa Mori, Masahiko Hunada, Miwa Misawa, and Hiroshi Nagase. Attenuation of the discriminative Stimulus Properties of Cocaine by δ -Opioid Receptor Antagonists. *European J. Pharmacol.*, 263, 207-211, **1994**.
- (63) Tstomu Suzuki, Michiharu Yoshiike, Hirokazu Mizoguchi, Junzo Kamei, Miwa Misawa, and Hiroshi Nagase. Blokade of δ -Opioid Receptors Prevents Morphine-Induced Place Preference in Mice. *Jpn. J. Pharmacol.*, 66, 131-137, **1994**.
- (62) Junzo Kamei, Yuriko Iwamoto, Miwa Misawa, Hiroshi Nagase, and Yutaka Kasuya. Antitussive Effect of (+) Pentazocine in Diabetic Mice Is Mediated by ρ -Sites, but Not by μ or κ -Opioid Receptor. *Jpn. J. Psychopharmacol.*, 14, 147-152, **1994**.
- (61) Junzo Kamei, Yuriko Iwamoto, Miwa Misawa, Hiroshi Nagase, and Yutaka Kasuya. Effects of Diabetes on the Antinociceptive Effect of (+)Pentazocine in Mice. *Res. Communication in Chem. Path. and Pharm.*, 84, 105-110, **1994**.
- (60) Junzo Kamei, Yuriko Iwamoto, Miwa Misawa, Hiroshi Nagase, and Yutaka Kasuya. Streptozocin- Induced Diabetes Selectively Enhances Aninociception Mediated by δ_1 -but not δ_2 -Opioid Receptors. *Life Sci.*, 55, 121-126, **1994**.
- (59) K. Arakawa, T. Akami, M. Okamoto, K. Akioka, P.C. Lee, Y. Sugano, J. Kamei, T. Suzuki, H. Nagase, Y. Tsutihashi, and T. Oka. Prolongation of Heart Xenograft Survival in the NK-Deficient Rat. *Transplantation Proceeding*, 26, 1266-1267, **1994**.
- (58) Junzo Kamei, Yuriko Iwamoto, Miwa Misawa, Hiroshi Nagase and Yutaka Kasuya. Antitussive Effect of L-arginine in Diabetic Mice. *European J. Pharmacol.*, 254, 113-117, **1994**.
- (57) Junzo Kamei, Yuriko Iwamoto, Miwa Misawa, Hiroshi Nagase and Yutaka Kasuya. Antitussive Effect of [Met⁵]enkephalin-Arg⁶-Phe⁷ in Mice. *European J. Pharmacol.*, 253, 293-296, **1994**.
- (56) Junzo Kamei, Yuriko Iwamoto, Tsutomu Suzuki, Miwa Misawa, Hiroshi Nagase and Yutaka Kasuya. Involvement of delta1 Opioid Receptor Antagonism in the Antitussive Effect of δ -Opioid Receptor Antagonists. *European J. Pharmacol.*, 251, 291-294, **1994**.
- (55) Junzo Kamei, Yuriko Iwamoto, Hideki Hitosugi, Miwa Misawa, Hiroshi Nagase and Yutaka Kasuya. Streptozocin-induced Diabetes Selectively Reduces Antinociception Mediated by μ_1 -Opioid Receptors, but not that Mediated by μ_2 -Opioid Receptors. *Neuroscience Lett.*, 165, 141-143, **1994**.

- (54) Junzo Kamei, Yuriko Iwamoto, Tsutomu Suzuki, Miwa Misawa, Yutaka Kasuya, Hiroshi Nagase, Takafumi Okutomi, Gen-ichiro Soma and Denichi Mizuno. Antinociceptive Effect of Lipopolysacchale from Pantoea Agglomerans on Streptozocin-Induced Diabetic Mice. *European J. Pharmacol.*, 251, 95-98, **1994**.
- (53) Junzo Kamei, Hideki Hitosugi, Miwa Misawa, Hiroshi Nagase and Yutaka Kasuya δ -Opioid Receptor-Mediated Forced Swimming Stress-induced Antinociception in the Formalin Test. *Pseuchopharmacol.*, 113, 15-18, **1993**.
- (52) Masahiko Funada, Tsutomu Suzuki, Minoru Narita, Miwa Misawa and Hiroshi Nagase. Blokade of Morphine Reward through the Activation of κ Opioid Receptors in Mice. *Neuropharmacol.*, 32, 1315-1323, **1993**.
- (51) Masahiko Funada, Tsutomu Suzuki, Minoru Narita, Misawa Miwa, and Hiroshi Nagase. Modification of Morphine-Induced Locomotor Activity by Pertussis Toxin: Biochemical and Behavioral Studies in Mice. *Brain Res.*, 619, 163-172, **1993**.
- (50) Junzo Kamei, Naoya Kawashima, Hideki Hitosugi, Miwa Misawa, Hiroshi Nagase, and Yutaka Kasuya. Effect of Diabetes on the antinociceptive Effect of β -Endorphin. *Brain Res.*, 619, 76-80, **1993**.
- (49) Junzo Kamei, Hideki Hitosugi, Miwa Misawa, Hiroshi Nagase, and Yutaka Kasuya. Cold Water Swim Stress Inhibits the Nociceptive response to intrathecally Administrated Somatostatin, but not Substance P. *Life Sci.*, 52, 169-174, **1993**.
- (48) Junzo Kamei, Yuriko Iwamoto, Miwa Misawa, Hiroshi Nagase, and Yutaka Kasuya. Antitussive Effect of β -Endorphine is Mediated by μ -Opioid Receptors, but not by κ - or ϵ -Opioid Receptors. *European J. Pharmacol.*, 233, 251-254, **1993**.
- (47) K. Arakawa, T. Akami, M. Okamoto, K. Akioka, I. Nakai, T. Oka and H. Nagase. Immuno-suppression by δ -Opioid Receptor Antagonist. *Transplantation Proceedings*, 25, 738-740, **1993**.
- (46) Junzo Kamei, Yuriko Iwamoto, Tsutomu Suzuki, Hiroshi Nagase, Miwa Misawa, and Yutaka Kasuya. Differential Modulation of μ -Opioid Receptor-Mediated Antitussive Activity by delta-Opioid Receptor Agonists in Mice. *Eur. J. Pharmacol.*, 234, 117-120, **1993**.
- (45) Junzo Kamei, Yuriko Iwamoto, Tsutomu Suzuki, Miwa Misawa, Hiroshi Nagase and Yutaka Kasuya. Antitussive Effects of Naltrindole, a Selective δ -Opioid Receptor Antagonist, in Mice and Rats. *Eur. J. Pharmacol.*, 249, 161-165, **1993**.
- (44) Junzo Kamei, Naoya Kawashima, Yuriko Iwamoto, Tsutomu Suzuki, Hiroshi Nagase, Miwa Misawa and Yutaka Kasuya. The Effect of Chronic Treatment with Naltrindole, a Selective delta Antagonist, on μ -Opioid Receptor-mediated

- Antinociception in Diabetic Mice. *Psychopharmacol.*, 113, 167-171, **1993**.
- (43) Minoru Narita, Tsutomu Suzuki, Masahiko Funada, Misawa Miwa, and Hiroshi Nagase. Involvement of δ -Opioid Receptors in the Effects of Morphine on Locomotor Activity and the Mesolimbic Dopaminergic System in Mice. *Psychopharmacology*, 111, 423-426, **1993**.
- (42) Minoru Narita, Kazuaki Takamori, Naoyu Kawashima, Masahiko Funada, Junzo Kamei, Tsutomu Suzuki, Misawa Miwa, and Hiroshi Nagase. Activation of Central ATP-Sensitive Potassium Channels Produces the Antinociception and Spinal Noradrenaline Turnover-Enhancing Effect in Mice. *Psychopharmacol.* 113, 11-14, **1993**.
- (41) Minoru Narita, Yuki Takahashi, Kazuaki Takamori, Masahiko Funada, Tsutomu Suzuki, Misawa Miwa, and Hiroshi Nagase. Effect of κ -Agonist on the Antinociception and Locomotor Enhancing Action Induced by Morphine in Mice. *Jpn.J. Pharmacol.*, 62, 15-24, **1993**.
- (40) Minoru Narita, Tsutomu Suzuki, Masahiko Funada, Misawa Miwa and Hiroshi Nagase. Blockade of the Morphine-induced increase in Turnover of Dopamine on the Mesolimbic Dopaminergic System by κ -Opioid Receptor Activation in mice. *Life Sciences*, 52, 397-404, **1993**.
- (39) Minoru Narita, Yuki Takahashi, Tsutomu Suzuki, Misawa Miwa and Hiroshi Nagase. An ATP-sensitive Potassium Channel Blocker Abolishes the Potentiating Effect of Morphine on the Bicuculline Induced Convulsion in mice. *Psychopharmacology*, 110, 500-502, **1993**.
- (38) Minoru Narita, Tsutomu Suzuki, Misawa Miwa and Hiroshi Nagase. Antagonism of the Morphine-Induced Straub Tail Reaction by κ -Opioid Receptor Activation in mice. *Psychopharmacology*, 110, 254-256, **1993**.
- (37) Junzo Kamei, Yukiko Iwamoto, Naoya Kawashima, Tsutomu Suzuki, Hiroshi Nagase, Miwa Misawa, and Yutaka Kasuya. Possible Involvement of μ_2 -Mediated Antitussive Activity in the Mouse. *Neurosci. Lett.*, 149, 169-172, **1993**.
- (36) Tsutomu Suzuki, Masahiko Funada, Minoru Narita, and Hiroshi Nagase. Morphine Induced Place Preference in the CXBK Mouse: Characteristics of μ Opioid Receptor Subtypes. *Brain Res.*, 602, 45-52, **1993**.
- (35) P. S. Portoghese, M. Sultana, H. Nagase and A.E. Takemori. A Highly Selective δ_1 -Opioid Receptor Antagonist: 7-Benzylidene Naltrexone. *European J. Pharmacol.*, 218, 195-196, **1992**.
- (34) Minoru Narita, Tsutomu Suzuki, Miwa Misawa, Hiroshi Nagase, Akemi Nabeshima, Takeshi Ashizawa, Hiroki Ozawa, Toshikazu Saito, and Naohiko

- Takahata. Role of Central ATP-sensitive Potassium Channels in the Analgesic Effect and Spinal Noradrenaline Turnover-Enhancing Effect Intracerebroventricularly Injected Morphine in Mice. *Brain Res.*, *596*, 209-214, **1992**.
- (33) Minoru Narita, Tsutomu Suzuki, Miwa Misawa and Hiroshi Nagase. Role of Central ATP-sensitive Potassium Channels in the Hyperthermic Effect of Morphine in mice. *Psychopharmacol.*, *109*, 239-240, **1992**.
- (32) K. Arakawa, T. Akami, M. Okamoto, H. Nakajima, M. Mitsuo, I. Nakai, T. Oka, Hiroshi Nagase and S. Matsutomo. Immunosuppressive Effect of δ -Opioid Receptor Antagonist on Xenogeneic Mixed Lymphocyte Reaction. *Transplantation Proceedings*, *24*, 696-697, **1992**.
- (31) Tsutomu Suzuki, Yoshinao Shiozaki, Yoshikazu Matsukawa, Miwa Misawa, and Hiroshi Nagase. Role of μ - and κ - Opioid Receptors in Cocaine-Induced Conditioned Place Preference. *Jpn. Pharmacol.*, *58*, 435-442, **1992**.
- (30) A. E. Takemori, M. Sultana, H. Nagase, and P. S. Portoghese. Agonist and Antagonist Activities of Ligands Derived from Naltrexone and Oxymorphone. *Life Sciences*, *50*, 1491-1495, **1992**.
- (29) T. Endoh, H. Matsuura, C. Tanaka, and H. Nagase. Nor-Binaltorphimine: A Potent and Selective kappa Opioid Receptor Antagonist with Long-Lasting Activity in vivo. *Arch. Int. Pharmacodyn. Ther.*, *316*, 30-42, **1992**.
- (28) Hisao Nishiyama, Kojun Isaoka, Kiyotaka Ohno, Hiroshi Nagase, Kazuhisa Matsumoto and Hideo Yoshiwara. Metal-Halogen Exchange between Polybromoanisoles and Aliphatic Grignard Reagents: A Synthesis of Cyclopenta[b]benzofurane. *J. Org. Chem.*, *50*, 407-410, **1992**.
- (27) Kohei Arakawa, Toshikazu Akami, Masahiko Okamoto, Takahiro Oka, and Hiroshi Nagase. The Immunosuppressive Effect of δ -Opioid Receptor Antagonist on Rat Renal Allograft Survival. *Transplantation*, *53*, No4, 951-953, **1992**.
- (26) Tsutomu Suzuki, Minoru Narita, Yuki Takahashi, Miwa Misawa, and Hiroshi Nagase. Effects of nor-Binaltorphimine on the Development of Analgesic Tolerance to and Physical Dependence on Morphine. *Eur. J. Pharmacol.*, *213*, 91-97, **1992**.
- (25) Tsutomu Suzuki, Masahiko Funada, Minoru Narita, Miwa Misawa, and Hiroshi Nagase. Pertussis toxin abolishes μ - and δ -Opioid Agonist-Induced Place Preference. *Eur. J. Pharmacol.*, *205*, 85-88, **1991**.
- (24) P. S. Portoghese, Hiroshi Nagase, K. E. Maloney-Huss, C.-E. Lin and A. E. Takemori. Role of Spacer and Address Components in Peptidomimetic δ Opioid

- Receptor Antagonists Related to Naltrindole. *J. Med. Chem.*, *34*, 1715-1720, **1991**.
- (23) Tsutomu Suzuki, Masahiko Funada, Minoru Narita, Miwa Misawa, and Hiroshi Nagase. Penazocine Induced Biphasic Analgesia in mice. *Life Sciences*, *48*, 1827-1835, **1991**.
- (22) P. S. Portoghese, A. Garzon-Aburgeh, Hiroshi Nagase, C-E. Lin, and A. E. Takemori. Role of the Spacer in Confering kappa Opioid Receptor Selectivity to Bivalent Ligands Related to Nornaltrorphimine. *J. Med. Chem.*, *34*, 1292-1296, **1991**.
- (21) Hiroshi Nagase, Kazuhisa Matsumoto, Hideo Yoshiwara, Atsuko Tajima, and Kiyotaka Ohno. Synthesis of (+)-5, 6, 7-Trinor-4, 8-Inter-*m*-Phenylene PGI₂. *Tetrahedron Lett.*, *31*, 4493-4494, **1990**.
- (20) Kiyotaka Ohno, Hisao Nishiyama, Hiroshi Nagase, Kazuhisa Matsumoto, and Mamoru Ishikawa. Synthesis of (±)-5, 6, 7-Trinor-4, 8-Inter-*m*-Phenylene PGI₂. *Tetrahedron Lett.*, *31*, 4489-4492, **1990**.
- (19) Tsutomu Suzuki, Yukiji Fukagawa, Miwa Misawa, and Hiroshi Nagase. "Paradoxical" Analgesia and Aggravated Morphine Dependence Induced by Opioid Antagonists. *Life Sciences*, *47*, 515-521, **1990**.
- (18) Hiroshi Nagase and P. S. Portoghese. Facile Intramolecular O-14-C-7 Acetyl Transfer in Opiate 14-Acetate Esters. *J. Org. Chem.*, *55*, 365-367, **1990**.
- (17) Hiroshi Nagase, Akira Abe, and P. S. Portoghese. The Facility of Formation of a 6-Δ-Bond in Dihydromorphinone and Related Opiates. *J. Org. Chem.*, *54*, 4120-4125, **1989**.
- (16) P. S. Portoghese, H. Nagase, and A. E. Takemori. Only One Pharmacophore Is Required for the κ Opioid Antagonist Selectivity of Norbinaltorphimine. *J. Med. Chem.*, *31*, 1344-1347, **1988**.
- (15) P. S. Portoghese, H. Nagase, A. W. Lipkowski, D. L. Larson, and A. E. Takemori. Binaltorphimine-Related Bivalent Ligands and Their kappa Opioid Receptor Antagonist Selectivity. *J. Med. Chem.*, *31*, 281-282, **1988**.
- (14) P. S. Portoghese, M. Sultana, H. Nagase, and A. E. Takemori. Application of Message-Address Concept in the Design of Highly Potent and Selective Non-Peptide δ-Opioid Receptor Antagonists. *J. Med. Chem.*, *31*, 836-841, **1988**.
- (13) A. W. Lipkowski, H. Nagase, and P. S. Portoghese. A Novel Pyrrole Synthesis via Reaction of Ketones with N-Aminoimide. *Tetrahedron Lett.*, 4257-4260, **1986**.
- (12) Haruki Niwa, Kazumasa Wakamatsu, Tsuneaki Hida, Kenji Niiyama Hideo Kigoshi, Mayumi Yamada, Hiroshi Nagase, Masaaki Suzuki, and Kiyoyuki

- Yamada. Stereocontrolled Total Synthesis of (-)-Picrotoxin and (+)-Coriamyrtin via a Common Isotwistane Intermediate. *J. Am. Chem. Soc.*, *106*, 4547-4552, **1984**.
- (11) Minoru Suzuki, Ikuo Morita, Mieko Kawamura, Sei-itsu Murata, Masato Nishizawa, Tadashi Miyake, Hiroshi Nagase, Kiyotaka Ohno, and Hirohiko Shimizu. Quantitative Determination of 6-keto Prostaglandin F₁α in Biological Fluids by Capillary Gas Chromatography-Chemical Ionization Mass Spectrometry. *J. Chromato.*, *221*, 361-366, **1980**.
- (10) Hisao Nishiyama, Hiroshi Nagase, and Kiyotaka Ohno. A Mild Methylation of Alcohols with Diazomethane Catalyzed by Silica Gel. *Tetrahedron Lett.*, 4671-4674, **1979**.
- (9) Kiyotaka Ohno, Hisao Nishiyama, and Hiroshi Nagase. Reaction of Some Amides and Thioamides with Diazomethane Catalyzed by Silica Gel. *Tetrahedron Lett.*, 4405-4406, **1979**.
- (8) Kiyoyuki Yamada, Shunsuke Gotoh, Hiroshi Nagase, Yoshinori Kyoutani, and Yoshimasa Hirata. Convenient Methods for Deoxygenation of Epoxides to Olefins. *J. Org. Chem.*, *43*, 2076-2077, **1978**.
- (7) Kiyoyuki Yamada, Shunsuke Gotoh, and Hiroshi Nagase. Total Synthesis and Stereostructure of (+)-Solavetivone, a Stress Metabolite from Infected Potato Tubers; X-Ray Crystal and Molecular Structure of an Intermediate in the Synthesis. *J. C. S. Chem. Comm.*, *344*, 554-555, **1977**.
- (6) Kiyoyuki Yamada, Kaneyoshi Katoh, Hiroshi Nagase, and Yoshimasa Hirata. Protection of Tertiary Hydroxyl Group as Methylthiomethyl Ethers. *Tetrahedron Lett.*, 65-66, **1976**.
- (5) Masaaki Suzuki, Yoshihiro Hayakawa, Kohzoh Aoki, Hiroshi Nagase, Hitoshi Nakamura, Kiyoyuki Yamada and Yoshimasa Hirata. Stereochemistry of Intermediates in the Syntheses of Dendrobium Alkaloids. *Tetrahedron Lett.*, 331-334, **1973**.
- (4) Kiyoyuki Yamada, Hiroshi Nagase, Yoshihiro Hayakawa, Kohzo Aoki, and Yoshimasa Hirata. Synthetic Studies on Spirovetivanes. II. Stereospecific Total Synthesis of *dl*-Hinesol, *dl*-α-Vetispirene, and *dl*-β-Vetispirene (*dl*-β-Isospirovetivone). *Tetrahedron Lett.*, 4967-4970, **1973**.
- (3) Kiyoyuki Yamada, Hiroshi Nagase, Yoshihiro Hayakawa, Kohzo Aoki, and Yoshimasa Hirata. Synthetic Studies on Spirovetivanes. I. Spirocondensation of a 4-(3'-Formylpropyl)-3-Cyclohexenone and Stereospecific Total Synthesis of *dl*-β-Vetivone. *Tetrahedron Lett.*, 4963-4966, **1973**.

- (2) Kiyoyuki Yamada, Masaaki Suzuki, Yoshihiro Hayakawa, Kohzo Aoki, Hiroshi Nagase, and Yoshimasa Hirata. Total Synthesis of *dl*-Dendrobine. *J. Amer. Chem. Soc.*, *94*, 8278-8280, **1972**.
- (1) Kiyoyuki Yamada, Matsuhiko Aratani, Yoshihiro Hayakawa, Hitoshi Nakamura, Hiroshi Nagase, and Yoshimasa Hirata. Facile Synthesis of Tricyclo[5.3.1.0.3.8]undecane and Spiro[5.5]undecane Systems from a Common Intermediate. *J. Org. Chem.*, *36*, 3653-3656, **1971**.

Reviews

- (51) 長瀬 博
「低分子オレキシン作動薬の創出ーナルコレプシー治療薬開発に向けて」
医学の歩み, *259*, 2 196-197, **2016**.
- (50) Hiroshi Nagase, Noriki Kutsumura
Synthesis of Novel Triplets with a 1,3,5-trioxazatriquinane Skeleton and their Pharmacologies for Opioid Receptors
The Archiv der Pharmazie, *348*, 375-389, **2015**.
- (49) 長瀬 博
「運と協力者に恵まれた創薬人生」
新規素材探索研究会会報, *12*, 2, 2014.
- (48) 長瀬 博
「難治性そう痒症治療薬ナルフラフィン塩酸塩の創出」「輝く 日本人による発見と新規開発」
医学の歩み, *251*, *11*, 1091-1094, **2014**.
- (47) 長瀬 博
難治性掻痒症治療薬レミッチ®カプセルの創出」
医学の門, *54*, *6*, **2014**.
- (46) 長瀬 博
難治性掻痒症治療薬ナルフラフィンの創出と痒み調節機構の解明
ペインクリニック, *35*, *11*, 1519-1526, **2014**.
- (45) 長瀬 博
「熾烈な新薬開発競争を振り返って ～κ受容体作動薬（モルヒネから依存性分離した薬物）はどう創製されたか～」
第58回日本透析医学会 学術集会・総会, 4-5, **2014**.
- (44) Hideaki Fujii, Toshihiro Takahashi, Hiroshi Nagase
Non-peptidic δ opioid receptor agonists and atagonists (2000-2012),

- Expert Opin. Ther. Patent* 23, 9, 1181-1208, **2013**.
- (43) Hiroshi Nagase and Hideaki Fujii
Essential Structure of the κ Opioid Receptor Agonist Nalfurafine for Binding to the κ Receptor
Current Pharmaceutical Design 19, 7400-7414, **2013**.
- (42) 長瀬 博
「オピオイドと痒み」
J. Visual Dermatol, 11: 714-717, **2012**.
- (41) Hideaki Fujii, Shigeto Hirayama and Hiroshi Nagase
Opioid Kappa Receptor Selective Agonist TRK-820 (Nalfurafine Hydrochloride) Pharmacology
ISBN:978-953-51-0222-9, 81-98, **2012**.
- (40) Hiroshi Nagase and Hideaki Fujii
Design and Synthesis of Novel Opioid Ligands and Their Pharmacologies
Heterocycles, 85, 8, 1821-1867, **2012**.
- (39) 長瀬 博
「産学のコラボレーションを含めた創薬の展望」オピオイド δ 作動薬の研究開発を題材に
MEDCHEM NEWS, 22, 3, 38-45, **2012**.
- (38) 藤井秀明, 平山重人, 長瀬 博
「受容体オリゴマーの新展開 -新たな創薬標的としての可能性」
化学, 66, 9, 70-71, **2011**.
- (37) 長瀬 博
世界初の依存性のないオピオイド κ 受容体作動薬の創出、座談会、世界初の選択的 κ 受容体作動性経口そう痒症改善剤レミッチの誕生秘話と今後の展望
鳥居薬品株式会社, 5-6, **2011**.
- (36) 長瀬 博
「新薬創製への道」(6) 難治性掻痒症治療薬ナルフラフィン塩酸塩の開発
Pharm.Tec Japan, 27, 13, 61-66, **2011**.
- (35) 長瀬 博
ナルフラフィン塩酸塩 (レミッチカプセル) の設計・合成とその薬理学的特性
日本緩和医療薬学雑誌, 3, 4, 65-76, 115-122, **2010**.
- (34) 長瀬 博, 川村 邦昭, 河合 孝治, 早川 潤
新規メカニズムに基づく難治性そう痒症治療薬ナルフラフィン塩酸塩の創出
有機合成化学協会誌, 68, 12, 1261-1271, **2010**.
- (33) 長瀬 博

- 依存性分離と戦い
ファルマシア、*46, 9*, 839-843, **2010**.
- (32) 内海 潤、長瀬 博
世界初の止痒薬 Nalfurafine の開発と意義—痒み知覚におけるオピオイドシステム
の新機構
バイオサイエンスとインダストリー、*68, 1*, 28-33, **2010**.
- (31) 長瀬 博、河合 孝治
新規メカニズムに基づく難治性そう痒症治療薬ナルフラフィンの創出
ファルマシア、*46, 6*, 521-525, **2010**.
- (30) 長瀬 博
新しい κ オピオイドアゴニスト
生体の科学・伝達物質と受容体、*60, 5*, 452-453, **2009**.
- (29) 長瀬 博
新しい NTI 誘導体
生体の科学・伝達物質と受容体、*60, 5*, 450-451, **2009**.
- (28) 長瀬 博
オピオイド κ 受容体作動薬 (TRK-820:ナルフラフィン塩酸塩) の止痒薬としての
開発
真興交易 (株) 医薬出版部、ペインクリニック、*30, 6*, 833-836, **2009**.
- (27) 長瀬 博
 δ および κ -オピオイド受容体作動薬の設計、合成とその薬理作用
Anesthesia 21 Century 10 83-32): 19-30 **2008**.
- (26) 長瀬 博
(巻頭言) 創薬研究と有機化学にかける思い
有機合成化学協会誌、*66, 3*, 193, **2008**.
- (25) 長瀬 博
(巻頭言) 大学における創薬研究・教育のあり方を
ファーマビジョンニュース、*10*, **2007**.
- (24) 大田 雅照、小尾 紀行、馬場 厚生、本多 利雄
(座談会) 創薬の魅力を伝える
ファルマシア、*43, 4*, 301-306, **2007**.
- (23) H. Fujii, and H. Nagase
Rational Drug Design of Selective Opioid Receptor antagonist TAN-821 and
Antagonist TAN-1014
Current Medicinal Chemistry, 13, 1109-1118, **2006**.
- (22) 長瀬 博、藤井 秀明

δ オピオイド受容体作動薬、TAN-67 の設計合成

有機合成化学協会誌, 64, 4, 2006.

- (21) 長瀬 博
ヒット化合物から開発化合物まで一止痒薬の開発を例として—
日本化粧品学会誌, 30, 3, 148-151, 2006.
- (20) 長瀬 博
δ 作用薬 (—) TAN-67 の薬理作用と光学異性体 (+) TAN-67 の発痛作用及び神経因性疼痛モデルへの適用
Medchem News, 12, 1, 10-12, 2002.
- (19) 溝口 広一、桜田 忍、長瀬 博
今後臨床応用が期待される開発中のオピオイドおよびその関連薬—デルモルフィン誘導体
緩和医療学, 4, 2, 128-134, 2002.
- (18) 田中 利明、富樫 裕子、梅内 秀郎、安藤 直生、川村 邦昭、岡野 清、遠藤 孝、亀井 淳三、長瀬 博
オピオイド κ 受容体作動薬 TRK-820 のマウスにおける止痒作用、臨床薬理
JPN J Clin Pharmacol Ther 31, 2, 2000.
- (17) 矢島 義識、長瀬 博
μ オピオイドジュヨウタイのspraysバリエーションによるサブタイプの存在
日薬理誌, 115, 315, 2000.
- (16) 藤田 智、中江 祐理、土田 英昭、長瀬 博、遠藤 孝、並木 昭義
Deltorphin (Opioid δagonist) が Guinea pig 摘出灌流心臓に及ぼす影響
循環制御, 19, 2, 1998.
- (15) 長瀬 博
薬の解説、非ペプチド性 δ オピオイド作用薬の合理的設計と合成
ファルマシア, 33, 8, 876-878, 1997.
- (14) 西尾 伸太郎、長瀬 博、加納 寛二、青木 茂、神林 義憲
安定型 PGI₂ 誘導体ベラプロストナトイウムの開発
薬学雑誌, 117, 8, 509-521, 1997.
- (13) 西尾 伸太郎、長瀬 博、加納 寛二、青木 茂、神林 義憲
安定 PGI₂ 誘導体ベラプロストナトリウムの開発
月刊薬事, 39, 7, 135-136, 1997.
- (12) 長瀬 博. 独創的新薬の合理的設計
日薬理誌, 110, 161-162, 1997.
- (11) 長瀬 博、松本 和久、西山 久男
新規な安定 PGI₂ 誘導体 m-フェニレン PGI₂ の合成研究—ベラプロストア

- トリウムの製造法開発
有機合成化学協会誌、54, 12, 47-58, 1996.
- (10) 長瀬 博
神経伝達物質と免疫 (I I) (オピオイド受容体)
ケミカルタイムス、8-13, 1995.
- (9) 長瀬 博
企業の研究室を訪ねる、東レ (株) 基礎研究所、独創的な研究に打ち込める土壌作りを
化学、49, 1994.
- (8) 長瀬 博
神経伝達物質と免疫 (I) (病は気から)
ケミカルタイムス、152-158, 1994.
- (7) 荒川 幸平、赤見 敏一、岡本 雅彦、岡 隆宏、長瀬 博、松本 修
NTI (δ オピオイド受容体アンタゴニスト) によるラット移植腎における免疫抑制効果
今日の移植、日本医学館、5, 2, 145-149, 1992.
- (6) Hiroshi Nagase and P.S. Portoghese
Abnormal Reaction of 4,5-Epoxydihydromorphinons and its Application to the Synthesis of Opioid Receptor Antagonists, J. Pharmacobio-Dyn, 15, s-41 (1990), Proceedings of the 17th Symposium on Progress in Organic Reaction and Syntheses, November 7, 8, 1991, Fukuoka.
- (5) Hiroshi Nagase, Minoru Narita, Tsutomu Suzuki, and Miwa Misawa
Supraspinal Analgesic Action of Pentazocine in Mice, The International Narcotic Research Conference (INRC) 1989, Page 465-468, Alan R. Liss, Inc. 1990.
- (4) Hiroshi Nagase and P.s. Portoghese
Synthesis of delta Opioid Antagonista and Their Structure Activity Relationships, J. Pharmacobio-Dyn, 13, s-132 (1990), Proceedings of the 17th Symposium on Pharmacological Activity and Mechanism, November 9, 10, 1987, Narashino.
- (3) Yukiji Fukagawa, Tsutomu Suzuki, Miwa Misawa, and Hiroshi Nagase
Effect of Chronic Treatment with Specific Antagonists on Analgesia and Physical Dependence on Morphine in Rats, The International Narcotics Research Conference (INRC) 1989, page 527-530, Alan R. Liss, Inc. 1990.
- (2) 長瀬 博、河合 孝治
4, 5 -エポキシジヒドロモルフィノンの異常反応オピオイドリセプターアンタゴ

ニストの合成への応用

有機合成化学協会誌、47, 4, 374-383, 1989.

- (1) K. Ohno, H. Nagase, K. Matsumoto, H. Nishiyama, S. Nishio
Advances in prostaglandin, thromboxane, and leukotriene research, Vol.15,
Stereoselective synthesis of 5,6,7-trinor-4,8-inter-m-phenylene-PGI₂ Derivat
ives and their Inhibitory Activities to Human Platelet Aggregation.
Raven Press, New York, 279-281, 1985.

Books

- (29) 長瀬 博
新規メカニズムに基づくそう痒症治療薬ナルフラフィン塩酸塩(レミッチカプセル)
の創出、医学図書出版株式会社、*透析療法ネクスト XII*, 65-76, 2011.
- (28) Hiroshi Nagase and hideaki Fujii
Synthesis of Novel Basic Skeletons Derived from Naltrexone, Chemistry of
Opioid , Top. Curr. Chem., 299, 187-237, 2011, Springer
- (27) Hiroshi Nagase and hideaki Fujii
Opioid in Preclinical and Clinical Trials, Chemistry of Opioid , Top. Curr.
Chem., 299, 29-62, 2011. Springer.
- (26) Hiroshi Nagase
[Chemistry of Opioids] Top Curr Chem, Volume editor, 299 2011.
- (25) 長瀬 博
慢性疼痛における薬剤選定と治療薬開発、*技術情報協会*、310-312, 2010.
- (24) 長瀬 博 監訳、D.A.Williams, T.L. Lemke 編集
「改良型新薬創製のための薬科学、(第三巻)」、テクノミック、2008.
- (23) Hiroshi Nagase, Hideaki Fujii
Rational Drug Design of δ Opioid Receptor Agonist TAN-67 from δ Opioid
Receptor Antagonist NTI, Topics in Heterocycl Chem.(2007) 8: 99-125, Spr
ing-Verlag Berlin Heideberg.
- (22) 長瀬 博
創薬科学・医薬化学 (ベーシック薬学教科書シリーズ)、*化学同人*、3-27, 2007.
- (21) D.A.Williams, T.L.Le Lemke 編集、長瀬 博 監訳
「改良型新薬創製のための薬科学、(第二巻)」、テクノミック、2007.
- (20) C.G.Wermuth 編集、長瀬 博 監訳
「最新創薬化学、第二版 (下巻)」、テクノミック、2006.
- (19) D.A.Williams, T.L.L Lemke 編集、長瀬 博 監訳

- 「改良型新薬創製のための薬科学、(第一巻)」、テクノミック、**2005**.
- (18) H.S.Smith、井上 哲夫、鈴木 勉、高橋 秀則、長瀬 博、成田 年、益田 律子、輪島 善一郎 監修
「痛みの治療薬、その基礎から臨床まで」、エルゼビアジャパン、**2005**.
- (17) 長瀬 博
「ここまで進んだゲノム医科学と疾患研究」(実験医学増刊) 最新のメディシナルケミストリーードラックライクな化合物ライブラリーの利用、*羊土社*、210-216, **2005**.
- (16) Jun Utsumi, Yuko Togashi, Hideo Umeuchi, Kiyoshi Okano, Toshiaki Tanaka, and Hiroshi Nagase
Antipururitic Activity of a Novel κ -Opioid Receptor Agonist, TRK-820, *IT CH, Marcel Dekker*, 107-114, **2004**.
- (15) C.G.Wermuth 編集、長瀬 博 監訳
「最新創薬化学、第二版(上巻)」、テクノミック、**2004**.
- (14) 長瀬 博
「鎮痛・オピオイド研究最前線」(鎮痛薬・オピオイドペプチド研究会編) 新規オピオイド κ 受容体作動薬 TRK-820 の止痒作用、*ミクス*、62-69. **2002**.
- (13) Gareth Thomas 著、長瀬 博 監訳
「製薬科学、メディシナルケミストのためのエッセンス」、テクノミック、東京、**2000**.
- (12) 長瀬 博、上野 新也、亀井 淳三
「オピオイド治療. 課題と新潮流」(鎮痛薬・オピオイドペプチド研究会編) オピオイドの持つ非鎮痛薬理作用(2) 新規 δ オピオイド受容体拮抗薬、TRK-851 の合理的設計とその鎮咳作用、*ミクス*、141-148, **2001**.
- (11) 長瀬 博、山本 尚
「創薬・薬物分子設計のコツ」、エルゼビヤ・サイエンスミクス、**2001**.
- (10) 長瀬 博、遠藤 孝、田中 利明
「オピオイドの基礎と臨床」(鎮痛薬・オピオイドペプチド研究会編) μ 、 δ および κ 受容体に関する研究(2) 新規 κ 受容体作動薬の薬理学的特徴、*ミクス*、144-154, **2000**.
- (9) 長瀬 博、今村 芳文
「オピオイドのすべて」(鎮痛薬・オピオイドペプチド研究会編) 創薬動向の面から)、*ミクス*、213-231, **1999**.
- (8) C.G.Wermuth 編集、長瀬 博 監訳
「最新創薬化学(下巻)」、テクノミック、**1999**.
- (7) 遠藤 孝、長瀬 博

- 「疼痛治療の現状と展望、臨床および基礎の立場から」(鎮痛薬・オピオイドペプチド研究会編) 新規 κ 作動薬、TRK-820 の薬理作用、ミクス、137-151, 1998.
- (6) C.G.Wermuth 編集、長瀬 博 監訳
「最新創薬化学 (上巻)」、テクノミック、1998.
- (5) 長瀬 博
「痛みの神経科学」鎮痛薬開発の現状と展望、メジカルビュー社、212-221, 1997.
- (4) 長瀬 博
「オピオイド、適性使用と最近の進歩」(鎮痛薬・オピオイドペプチド研究会編)、オピオイド受容体選択性リガンドの合理的設計と合成、ミクス、167-181, 1997.
- (3) 野崎 正勝、長瀬 博
「創薬化学」、化学同人、1995.
- (2) 長瀬 博
「オピオイド、化学物質が解き明かす生体の謎」(化学増刊 120、鎮痛薬・オピオイドペプチド研究会編)、オピオイドリガンドの分子設計— κ および δ アンタゴニストの合成、化学同人、151-157, 1991.
- (1) Hiroshi Nagase
New Leads in Opioid Research, Proceedings of the international Narcotic Research Conference, Noordwijkerhout, The Netherlands, (Exc epta Medica), p 8-1, July 1990, Amsterdam-New York, 1990

Invited Lectures

- (53) 小野薬品工業株式会社
演題：オレキシン受容体リガンドの探索研究
日時：2016年11月2日
- (52) 厚木市皮膚科医会例会
演題：ナルフラフィンの研究開発と新規痒みの機序の解明
日時：2016年10月20日
- (51) 富山化学工業株式会社
演題：ナルフラフィンの研究開発、オレキシン受容体リガンドの探索研究、トリプレト薬の研究と化合物ライブラリーへの応用
日時：2016年10月14日
- (50) The 12th JSH Single Topic Conference
演題：Design and Synthesis of Nalfurafine for an Antipruritic Drug for Chronic liver Disease Patients and kidney Dialysis Patients

- 日時：2016年9月23日
- (49) Tsukuba Global Science Week 2016
演題：Design and Synthesis of Orexin 2 Receptor Agonists
日時：2016年9月19日
- (48) FIP2016
演題：The science and the development of non-addictive opioid receptor agonists
日時：2016年8月30日
- (47) 第36回鎮痛薬・オピオイドペプチンポジウム
演題：オレキシン2受容体作動薬の設計・合成
日時：2016年8月19日
- (46) Orexin Expert Forum
演題：医薬標的としてのオレキシン系
日時：2016年7月16日
- (45) 日本睡眠学会第41回定期学術集会
演題：オレキシン受容体作動薬：創薬への展望
日時：2016年7月7日
- (44) 第13回GPCR研究会
演題： κ 受容体拮抗薬ナルフラフィンの開発と痒みの発生機序
日時：2016年5月13日
- (43) 技術情報協会
演題：難治性掻痒症治療薬の研究開発と痒みの最新メカニズム
日時：2016年4月19日
- (42) 全薬工業株式会社
演題：オレキシン作動薬の設計・合成と薬理作用
日時：2016年3月11日
- (41) 脳・肝インターフェースメディシン研究センター シンポジウム
演題：オレキシン2受容体選択的作動薬の研究開発とその薬理作用
演題：Design and Synthesis of OX2R selective agonists and their pharmacologies
日時：2016年3月7日
- (40) 持田製薬株式会社創薬研究所
演題：オレキシン関連創薬
日時：2016年1月19日
- (39) Pacifichem 2015
演題：Design and synthesis of novel ligands of opioid receptors
日時：2015年12月17日

- (38) 第133回日本薬理学会関東部会ミニシンポジウム
演題：ナルフラフィンの OX1R 拮抗作用と U-50488H の嫌悪作用との関係
日時：2015年10月10日
- (37) 第18回近畿薬剤師学術大会 in 神戸
演題：患者ニーズに根ざした新薬開発 — 難治性そう痒症治療薬を例にして —
日時：2015年8月30日
- (36) テンプル大学
演題：Study and Development of Opioid κ Agonist, Nalfurafine and their Clinical Effects
日時：2015年7月27日
- (35) Drug Discovery & Therapy World Congress 2015
演題：DESIGN AND SYNTHESIS OF NOVEL δ OPIOID AGONISTS AND THEIR PHARMACOLOGIES
日時：2015年7月23日
- (34) 北海道大学
演題：難治性搔痒症治療薬ナルフラフィンの研究開発
日時：2015年7月16日
- (33) 大日本住友製薬株式会社
演題：難治性そう痒治療薬の研究開発
日時：2015年7月6日
- (32) 富士フィルム株式会社
演題：難治性そう痒治療薬の研究開発と痒みの最新メカニズム
日時：2015年6月30日
- (31) 株式会社 技術情報協会
演題：痒みの最新メカニズムと難治性搔痒症治療薬のポイント
日時：2015年3月31日
- (30) 第88回日本薬理学会年会
演題： δ 作動薬の設計・合成
日時：2015年3月20日
- (29) 脳肝インターフェースメディスンセンター 分子神経科学部門シンポジウム 2015
演題：難治性搔痒症治療薬ナルフラフィンの研究開発とその作用機序
日時：2015年3月13日
- (28) ライフイノベーション学位プログラム 第二回シンポジウム
演題：Study and development of opioid κ agonist, nalfurafine and their clinical effects
日時：2015年1月21日

- (27) DPhG Jahrestagung 2014
演題：Synthesis of a novel opioid receptor agonist, SYK-146 with 1,3,5-trioxazatriquinane skeleton and its pharmacologies
日時：2014年9月23日～26日
- (26) 都学研国際シンポジウム, 2014年9月4日～6日(東京)
演題：design and synthesis of novel ligands for opioid receptors
日時：2014年9月4日～6日
- (25) 協和発酵キリン
演題：ナルフラフィンの研究開発と最近の研究
日時：2014年8月22日
- (24) 鳥居薬品
演題：難治性掻痒症治療薬レミッチカプセルの創出
日時：2014年7月24日
- (23) R&D支援センター
演題：難治性掻痒症治療薬の研究開発と臨床試験のポイント
日時：2014年6月24日
- (22) University of Minnesota
演題：Design and Synthesis of Opioid Receptor Type Selective Ligands
日時：2014年6月
- (21) 大正製薬 医薬化学研究所
演題：オピオイド受容体タイプ選択的薬物の研究開発
日時：2014年6月6日
- (20) 金沢大学
演題：2つの独創的新薬(ドルナー、ナルフラフィン)の創製
日時：2014年5月22日
- (19) 東レ株式会社
演題：オピオイド選択的薬物の設計と合成
日時：2014年2月28日
- (18) 全薬工業株式会社 中央研究所
演題：創薬化学探索研究から開発まで
日時：2014年2月14日
- (17) 名古屋大学トランスフォーマティブ生命分子研究所 (ITbM)
演題：ベラプロスト、ナルフラフィンの開発とオレキシン作動薬の研究
日時：2014年2月5日
- (16) 住友化学 健康・農業関連事業研究所
演題：プロスタグランジンおよびオピオイド系医薬の創薬研究

- 日時：2014年1月31日
- (15) 有機合成化学協会関東支部ミニシンポジウム千葉2013
演題：オピオイド受容体タイプ選択的リガンドの設計・合成
日時：2013年11月18日
- (14) The 2nd HBP Symposium at Tsukuba Global Science Week 2013
演題：Synthesis of Novel Triplets with 1,3,5-Trioxazatriquinane Skeleton Using Nitrogen Clamp
日時：2013年10月2日
- (13) 第7回日本緩和医療薬学会年会
演題：2つの薬が世に出るまで：安定型プロスタグランジン I₂ 誘導体（ドルナー）および難治性掻痒症治療薬レミッチカプセルの探索研究から発売まで
日時：2013年9月15日
- (12) 国立がん研究センター研究所
演題：オピオイド受容体拮抗薬の原虫症治療薬としての効果およびその作用機序（癌治療薬としての可能性）
日時：2013年8月26日
- (11) 旭化成ファーマ株式会社
演題：ベラプロストとナルフラフィンの研究開発
日時：2013年7月26日
- (10) 田辺三菱研究所
演題：ドルナーとナルフラフィンの研究開発
日時：2013年7月4日
- (9) 日本透析医学会学術集会・総会
演題：熾烈な新薬開発競争を振り返って～□受容体作動薬（モルヒネから依存性を分離した薬物）はどう創製されたか～
日時：2013年6月21日
- (8) 東レ医薬研究所
演題：オピオイド受容体選択的薬物の設計と合成
日時：2013年3月19日
- (7) 全薬工業株式会社
κ オピオイド受容体作動薬ナルフラフィン塩酸塩の研究開発
日時：2013年3月8日
- (6) 筑波大学
分子行動科学研究コア医科学セミナー
演題：難治性掻痒症治療薬・ナルフラフィン塩酸塩の研究開発
日時：2012年11月21日

- (5) 東京大学弥生講堂 第17回分生研シンポジウムアカデミア創薬～分子標的と化合物の力～
演題：難治性咳嗽治療薬、TRK-851の研究開発
日時：2012年10月29日
- (4) New Trends in Pain Research from Basic Research to Clinical Translation
Invited Opening Lecture in Italy
演題：Translational Research in Itching: Research and Development of nalfurafine Hydrochloride and its Application to Kidney Dialysis Patients with intractable pruritus.
日時：2012年9月13日
- (3) 技術情報協会セミナー痒みのメカニズム/評価法と新薬開発戦略
演題：難治性掻痒症治療薬ナルフラフィン塩酸塩の研究開発
日時：2012年8月20日
- (2) 日本薬理学会関東支部会招待講演
演題：ナルフラフィン塩酸塩の研究開発
日時：2012年7月14日
- (1) Wuxi pharma Tech社（上海の合成/アッセイ・GLP対応試験受託会社）招待講演
演題：Design and Synthesis of a Novel Antipruritic Agent, Nalfurafine Hydrochloride.
日時：2012年6月4日

Patents

- (25) オレキシン1受容体拮抗薬
2015-212553
長瀬 博、山本 直司、入鹿山 容子、斉藤 毅
- (24) スルホンアミド誘導体またはその薬学的に許容される酸付加塩
2015-119785
長瀬 博、柳沢 正史、斉藤 毅、沓村 憲樹、入鹿山 容子
- (23) モルヒナン誘導体
特願 2015-54079
長瀬 博、藤井 秀明、斎藤 顕宜、中田 恵理子、廣瀬 雅朗、大井 勲、林田 康平
- (22) アミド誘導体またはその薬学的に許容される酸付加塩
2015-31041
柳沢正史、長瀬 博、入鹿山 容子、斉藤 毅

- (21) スルホンアミド誘導体またはその薬学的に許容される酸付加塩
PCT/JP2014/082961
長瀬 博、永原 崇志
- (20) オレキシン受容体拮抗剤
特願 2014-222969
長瀬 博、入鹿山 容子、小川 靖裕、宮本 美津子、新田 一功
- (19) ナルフラフィン含有局所適用製剤
特願 2014-201237
長瀬 博、佐伯 健、下山 潤、多田 麻希子
- (18) スルホンアミド誘導体またはその薬学的に許容される酸付加塩
特願 2013-257523
長瀬 博、永原崇志
- (17) プロペラン誘導体
特願 2013-66941
長瀬 博、藤井秀明
- (16) モルヒナン誘導体
特願 2013-47325
長瀬 博、藤井秀明、中田恵理子、渡辺義和、高橋俊弘
- (15) モルヒナン誘導体
特願 2013-47324
長瀬 博、藤井秀明、中田恵理子、渡辺義和、高橋俊弘
- (14) プロペラン誘導体
特願 2013-12104
長瀬 博、藤井秀明
- (13) モルヒナン誘導体
特願 2012-72868
長瀬 博、藤井秀明、中田恵理子、渡辺義和、高橋俊弘
- (12) モルヒナン誘導体
特願 2012-168402
長瀬 博、藤井秀明、中田恵理子、渡辺義和、高橋俊弘
- (11) モルヒナン誘導体
特願 2012-168401
長瀬 博、藤井秀明、中田恵理子、渡辺義一、高橋俊弘
- (10) モルヒナン誘導体
特願 2012-51741
中田恵理子、藤井秀明、長瀬 博

- (9) 掻痒症改善経皮吸収貼付類
特願 2012—9168
長瀬 博、有吉知加子、原 和好、丸山 喜通
- (8) ジフェニルメチルピペラジン誘導体およびそれを含有する医薬組成物
特許番号：第 4782441 号
竹内 勤、長瀬 博、長 由美子、小林 正規
- (7) 6,14-エポキシモルヒナン誘導体及びその医薬用途
特開 2009—51749
長谷部 光、中島 真弓、長瀬 博
- (6) オキサビシクロ「2. 2. 2」オクタンを骨格とするモルヒナン誘導体及びその医薬用途
特開 2009—196933
長谷部 光、中島 真弓、長瀬 博
- (5) 4a,9a-架橋-ヘキサヒドロ-1H-インデノピリジン誘導体、その医薬用途、及びその製造方法
特開 2008—179556
木綿 しのぶ、泉本 直樹、長瀬 博
- (4) ピロール縮合モルヒナン誘導体及びその医薬用途
特開 2008—179554
長谷部 光、木綿 しのぶ、長瀬 博
- (3) 薬剤耐性解除用の医薬
特開 2008—37790
長瀬 博、長 由美子、竹内 勤、小林 正規、狩野 繁之
- (2) オピオイドδ受容体アゴニスト
特願 2008—522628
長瀬 博 (国際出願 WO 2008/001859 A1) .
- (1) 止痒剤
特願 2002—311186
長瀬 博、内海 潤、遠藤 孝、田中 利明、川村 邦明